







Match-up database Analyses Report

SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM

TSG-LEGOS-DM

North Atlantic Ocean

prepared by the Pi-MEP Consortium

March 15, 2019

Contents

1 Overview						
2	The	MDB file datasets				
	2.1	Satellite SSS product				
		2.1.1 SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM				
	2.2	In situ SSS dataset				
	2.3	Auxiliary geophysical datasets				
		2.3.1 CMORPH				
		2.3.2 ASCAT				
		2.3.3 ISAS				
		2.3.4 World Ocean Atlas Climatology				
	2.4	Overview of the Match-ups generation method				
		2.4.1 In Situ/Satellite data filtering				
		2.4.2 In Situ/Satellite Co-localization				
		2.4.3 MDB pair Co-localization with auxiliary data and complementary infor-				
		mation				
		2.4.4 Content of the Match-Up NetCDF files				
	2.5	MDB characteristics for the particular in situ/satellite pairs				
	-	2.5.1 Number of paired SSS data as a function of time and distance to coast				
		2.5.2 Histograms of the SSS match-ups				
		2.5.3 Distribution of in situ SSS depth measurements				
		2.5.4 Spatial Distribution of Match-ups				
		2.5.5 Histograms of the spatial and temporal lags of the match-ups pairs				
	MID	D CL A 1				
3		B file Analyses				
	3.1	Spatial Maps of the Temporal mean and Std of in situ and satellite SSS and of				
	2.0	the difference (ΔSSS)				
	3.2	Time series of the monthly averaged mean and Std of in situ and satellite SSS and				
	9.9	of the (ΔSSS)				
	3.3	Zonally-averaged Time-mean and temporal Std of in situ and satellite SSS and of				
	9 4	the ΔSSS				
	3.4	Scatterplots of satellite vs in situ SSS by latitudinal bands				
	3.5	Time series of the monthly averaged mean and Std of the Δ SSS sorted by latitudical bands				
	9.6	dinal bands				
	3.6	ASSS sorted as function of geophysical parameters				
	3.7	ΔSSS maps and statistics for different geophysical conditions				
4	Sun	nmary				
L	ist	of Figures				
	1	Number of match-ups between TSG-LEGOS-DM and SMOS-L3-CATDS-CECOS-				
	1	LOCEAN-V3-18DAYS-25KM SSS as a function of time (a) and as function of the				
		distance to coast (b) over the North Atlantic Ocean Pi-MEP region and for the				
		full satellite product period				



2	Histograms of SSS from TSG-LEGOS-DM (a) and SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM (b) considering all match-up pairs per bins of 0.1 over the North Atlantic Ocean Pi-MEP region and for the full satellite product	
3	period	18
4	1°x1° boxes and for the full satellite product period (b)	19
5	period	19
6	CECOS-LOCEAN-V3-18DAYS-25KM SSS product	20
7	(Satellite - TSG-LEGOS-DM). Only match-up pairs are used to generate these maps	21
•	TSG-LEGOS-DM) and Std of Δ SSS (Satellite - TSG-LEGOS-DM) over the North Atlantic Ocean Pi-MEP region considering all match-ups collected by the Pi-MEP	
8	platform. Left panel: Zonally averaged time mean SSS from SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM (black) and from TSG-LEGOS-DM (blue). Right	22
	panel: zonally averaged time-mean Δ SSS (Satellite - TSG-LEGOS-DM) for all the collected Pi-MEP match-up pairs estimated over the full satellite product period.	23
9	Contour maps of the concentration of SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM SSS (y-axis) versus TSG-LEGOS-DM SSS (x-axis) at match-up pairs for different latitude bands. For each plot, the red line shows x=y. The	
	black thin and dashed lines indicate a linear fit through the data cloud and the $\pm 95\%$ confidence levels, respectively. The number match-up pairs n , the slope and R^2 coefficient of the linear fit, the root mean square (RMS) and the mean	
	bias between satellite and in situ data are indicated for each latitude band in each plots	24
10	Monthly-average mean (red curves) Δ SSS (Satellite - TSG-LEGOS-DM) and ± 1 Std (black vertical thick bars) as function of time for all the collected Pi-MEP match-up pairs estimated over the North Atlantic Ocean Pi-MEP region and for	
	the full satellite product period are shown for different latitude bands: (a) Latitude band 80°S-80°N, (b) latitude band 20°S-20°N, (c) Mid Latitude bands 40°S-20°S	25
11	and 20°N-40°N and (d) Latitude bands 60°S-40°S and 40°N-60°N	25
	d) and distance to coast (e). In all plots the mean and Std of Δ SSS for each bin is indicated by the red curves and black vertical thick bars (± 1 Std)	26

12	Temporal mean gridded over spatial boxes of size 1°x1° of Δ SSS (SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM - TSG-LEGOS-DM) for 5 different subdatasets corresponding to:RR=0 mm/h, $3 < U_{10} < 12$ m/s, SST>5°C, distance to coast > 800 km (a), RR=0 mm/h, $3 < U_{10} < 12$ m/s (b), RR>1mm/h and $U_{10} < 4$ m/s (c), WOA2013 SSS Std<0.2 (d), WOA2013 SSS Std>0.2 (e)	27	
13	Normalized histogram of Δ SSS (SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-	21	
10	25KM - TSG-LEGOS-DM) for 5 different subdatasets corresponding to: RR=0		
	mm/h , $3 < U_{10} < 12 \text{ m/s}$, $SST > 5$ °C, distance to coast $> 800 \text{ km}$ (a), $RR = 0 \text{ mm/h}$,		
	$3 < U_{10} < 12 \text{ m/s (b)}$, RR>1mm/h and $U_{10} < 4\text{m/s (c)}$, WOA2013 SSS Std<0.2		
	(d), WOA2013 SSS Std>0.2 (e)	28	



Acronym

Aquarius NASA/CONAE Salinity mission

ASCAT Advanced Scatterometer

ATBD Algorithm Theoretical Baseline Document

BLT Barrier Layer Thickness CMORPH CPC MORPHing technique

CTD Instrument used to measure the conductivity, temperature, and pressure of

seawater

DM Delayed Mode
EO Earth Observation
ESA European Space Agency
FTP File Transfer Protocol

GOSUD Global Ocean Surface Underway Data GTMBA The Global Tropical Moored Buoy Array

Ifremer Institut français de recherche pour l'exploitation de la mer

IPEV Institut polaire français Paul-Émile Victor

IQR Interquartile range ISAS In Situ Analysis System

Kurt Kurtosis (fourth central moment divided by fourth power of the standard de-

viation)

L2 Level 2

LEGOS Laboratoire d'Etudes en Géophysique et Océanographie Spatiales

LOCEAN Laboratoire d'Océanographie et du Climat : Expérimentations et Approches

Numériques

LOPS Laboratoire d'Océanographie Physique et Spatiale

MDB Match-up Data Base

MEOP Marine Mammals Exploring the Oceans Pole to Pole

MLD Mixed Layer Depth

NCEI National Centers for Environmental Information

NRT Near Real Time

NTAS Northwest Tropical Atlantic Station

OI Optimal interpolation

Pi-MEP Pilot Mission Exploitation Platform

PIRATA Prediction and Researched Moored Array in the Atlantic

QC Quality control

 R_{sat} Spatial resolution of the satellite SSS product

RAMA Research Moored Array for African-Asian-Australian Monsoon Analysis and

Prediction

r² Square of the Pearson correlation coefficient

RMS Root mean square

RR Rain rate

SAMOS Shipboard Automated Meteorological and Oceanographic System

Skew Skewness (third central moment divided by the cube of the standard deviation)

SMAP Soil Moisture Active Passive (NASA mission)
SMOS Soil Moisture and Ocean Salinity (ESA mission)
SPURS Salinity Processes in the Upper Ocean Regional Study

SSS Sea Surface Salinity

 SSS_{insitu} In situ SSS data considered for the match-up



 SSS_{SAT} Satellite SSS product considered for the match-up

 Δ SSS Difference between satellite and in situ SSS at colocalized point (Δ SSS =

 SSS_{SAT} - SSS_{insitu})

SST Sea Surface Temperature Std Standard deviation

Std* Robust Standard deviation = median(abs(x-median(x)))/0.67 (less affected by

outliers than Std)

Stratus Surface buoy located in the eastern tropical Pacific

Survostral SURVeillance de l'Océan AuSTRAL (Monitoring the Southern Ocean)

TAO Tropical Atmosphere Ocean

TSG ThermoSalinoGraph

WHOI Woods Hole Oceanographic Institution
WHOTS WHOI Hawaii Ocean Time-series Station

WOA World Ocean Atlas



1 Overview

In this report, we present systematic analyses of the Match-up DataBase (MDB) files generated by the Pi-MEP platform within the following Pi-MEP region and for the below pair of Satellite/In situ SSS data:

- Pi-MEP region: North Atlantic Ocean (download the corresponding mask here)
- SSS satellite product (SSS $_{SAT}$): SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM
- In situ dataset (SSS_{Insitu}): TSG-LEGOS-DM (download the corresponding report here)

In the following, $\Delta SSS = SSS_{SAT}$ - SSS_{Insitu} denotes the difference between the satellite and in situ SSS at the colocalized points that form the MDB.

This report presents successively:

The MDB file DataSets (Section 2)

- A short description of the satellite SSS product considered in the match-up (2.1)
- A short description of the In situ SSS dataset considered in the match-up (2.2)
- A short description of the auxiliary geophysical datasets co-localized with SSS pairs (2.3)
- An overview of how the Match-ups were evaluated (2.4)
- An overview of the MDB characteristics for the particular in situ/satellite pairs (2.5)

The major results of the MDB file Analyses (Section 3)

- Spatial Maps of the Time-mean and temporal Std of in situ and satellite SSS and of the Δ SSS (3.1)
- Time series of the monthly averaged mean and Std of in situ and satellite SSS and of the Δ SSS (3.2)
- Zonally-averaged Time-mean and temporal Std of in situ and satellite SSS and of the Δ SSS (3.3)
- Scatterplots of satellite vs in situ SSS by latitudinal bands (3.4)
- Time series of the monthly averaged mean and Std of the Δ SSS sorted by latitudinal bands (3.5)
- Δ SSS sorted as function of geophysical parameters (3.6)
- Δ SSS maps and statistics for different geophysical conditions (3.7)

All analyses are conducted over the Pi-MEP Region specified above and over the full satellite SSS product period.



2 The MDB file datasets

2.1 Satellite SSS product

2.1.1 SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM

In order to propose improved methodologies to be implemented in the near real time CATDS processing chain (CATDS-CPDC), LOCEAN/IPSL (UMR CNRS/UPMC/IRD/MNHN) and ACRIst have derived a methodology for correcting systematic SSS biases. The v3 release uses an improved 'debiasing' technique (Boutin et al. (2018)) and the adjustment of the long term mean SMOS SSS in very dynamical areas, like in river plumes, and the bias correction at high latitudes have been improved.

Maps are provided every 4 days from 01/2010 to 12/2017 and are derived from a combination of ascending and descending orbits. Debiased SSS are temporally averaged using a slipping Gaussian kernel with a full width at half maximum of 18 days (18 day product). Maps are at a spatial resolution 25x25km2 and a median filtering over nearest neighbors is applied.

The L3 DEBIAS LOCEAN v2 Sea Surface Salinity product (Boutin et al. (2017)) have been produced by LOCEAN/IPSL (UMR CNRS/UPMC/IRD/MNHN) laboratory and ACRI-st company and is distributed by the Ocean Salinity Expertise Center (CEC-OS) of the CNES-IFREMER Centre Aval de Traitement des Donnees SMOS (CATDS), at IFREMER, Plouzane (France).

SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-							
Spatial resolution	$25~\mathrm{km}$						
Temporal resolution	18 days (file every 4 days)						
Temporal coverage	From 2010-01-16 to 2017-12-26						
Data Provider	CATDS						
Release Date	2018-07						
Version	3						
Data access	$debiasedSSS_18days_v3$						
DOI	http://doi.org/10.17882/52804						
Documentation	Doc_L3_DEBIAS_LOCEAN_v3.pdf						

Table 1: Satellite SSS product characteristics

2.2 In situ SSS dataset

The TSG-LEGOS-DM dataset correspond to sea surface salinity delayed mode data derived from voluntary observing ships collected, validated, archived, and made freely available by the French Sea Surface Salinity Observation Service (Alory et al. (2015)). Adjusted values when available and only collected TSG data that exhibit quality flags=1 and 2 were used.

2.3 Auxiliary geophysical datasets

Additional EO datasets are used to characterize the geophysical conditions at the in situ/satellite SSS pair measurement locations and time, and 10 days prior the measurements to get an estimate of the geophysical condition and history. As discussed in Boutin et al. (2016), the presence of vertical gradients in, and horizontal variability of, sea surface salinity indeed complicates comparison of satellite and in situ measurements. The additional EO data are used here to get



a first estimates of conditions for which L-band satellite SSS measured in the first centimeters of the upper ocean within a 50-150 km diameter footprint might differ from pointwise in situ measurements performed in general between 10 and 5 m depth below the surface. The spatiotemporal variability of SSS within a satellite footprint (50–150 km) is a major issue for satellite SSS validation in the vicinity of river plumes, frontal zones, and significant precipitation. Rainfall can in some cases produce vertical salinity gradients exceeding 1 pss m⁻¹; consequently, it is recommended that satellite and in situ SSS measurements less than 3-6 h after rain events should be considered with care when used in satellite calibration/validation analyses. To identify such situation, the Pi-MEP test platform is first using CMORPH products to characterize the local value and history of rain rate and ASCAT gridded data are used to characterize the local surface wind speed and history. For validation purpose, the ISAS monthly SSS in situ analysed fields at 5 m depth are collocated and compared with the satellite SSS products. The use of ISAS is motivated by the fact that it is used in the SMOS L2 official validation protocol in which systematic comparisons of SMOS L2 retrieved SSS with ISAS are done. In complement to ISAS, monthly std climatological fields from the World Ocean Atlas (WOA13) at the match-up pairs location and date are also used to have an a priori information of the local SSS variability.

2.3.1 CMORPH

Precipitation are estimated using the CMORPH 3-hourly products at 1/4° resolution (Joyce et al. (2004)). CMORPH (CPC MORPHing technique) produces global precipitation analyses at very high spatial and temporal resolution. This technique uses precipitation estimates that have been derived from low orbiter satellite microwave observations exclusively, and whose features are transported via spatial propagation information that is obtained entirely from geostationary satellite IR data. At present NOAA incorporate precipitation estimates derived from the passive microwaves aboard the DMSP 13, 14 and 15 (SSM/I), the NOAA-15, 16, 17 and 18 (AMSU-B), and AMSR-E and TMI aboard NASA's Aqua, TRMM and GPM spacecraft, respectively. These estimates are generated by algorithms of Ferraro (1997) for SSM/I, Ferraro et al. (2000) for AMSU-B and Kummerow et al. (2001) for TMI. Note that this technique is not a precipitation estimation algorithm but a means by which estimates from existing microwave rainfall algorithms can be combined. Therefore, this method is extremely flexible such that any precipitation estimates from any microwave satellite source can be incorporated.

With regard to spatial resolution, although the precipitation estimates are available on a grid with a spacing of 8 km (at the equator), the resolution of the individual satellite-derived estimates is coarser than that - more on the order of 12×15 km or so. The finer "resolution" is obtained via interpolation.

In effect, IR data are used as a means to transport the microwave-derived precipitation features during periods when microwave data are not available at a location. Propagation vector matrices are produced by computing spatial lag correlations on successive images of geostationary satellite IR which are then used to propagate the microwave derived precipitation estimates. This process governs the movement of the precipitation features only. At a given location, the shape and intensity of the precipitation features in the intervening half hour periods between microwave scans are determined by performing a time-weighting interpolation between microwave-derived features that have been propagated forward in time from the previous microwave observation and those that have been propagated backward in time from the following microwave scan. NOAA refer to this latter step as "morphing" of the features.

For the present Pi-MEP products, we only considered the 3-hourly products at 1/4 degree resolution. The entire CMORPH record (December 2002-present) for 3-hourly, 1/4 degree lat/lon resolution can be found at: ftp://ftp.cpc.ncep.noaa.gov/precip/CMORPH_V1.



O/RAW/. CMORPH estimates cover a global belt (-180°W to 180°E) extending from 60°S to 60°N latitude and are available for the complete period of the Pi-MEP core datasets (Jan 2010-now).

2.3.2 ASCAT

Advanced SCATterometer (ASCAT) daily data produced and made available at Ifremer/CERSAT on a 0.25°x0.25° resolution grid (Bentamy and Fillon (2012)) since March 2007 are used to characterize the mean daily wind at the match-up pair location as well as the wind history during the 10-days period preceding the in situ measurement date. These wind fields are calculated based on a geostatistical method with external drift. Remotely sensed data from ASCAT are considered as observations while those from numerical model analysis (ECMWF) are associated with the external drift. The spatial and temporal structure functions for wind speed, zonal and meridional wind components are estimated from ASCAT retrievals. Furthermore, the new procedure includes a temporal interpolation of the retrievals based on the complex empirical orthogonal function (CEOF) approach, in order to enhance the sampling length of the scatterometer observations. The resulting daily wind fields involves the main known surface wind patterns as well as some variation modes associated with temporal and spatial moving features. The accuracy of the gridded winds was investigated through comparisons with moored buoy data in Bentamy et al. (2012) and resulted in rms differences for wind speed and direction are about 1.50 m.s⁻¹ and 20°.

2.3.3 ISAS

The In Situ Analysis System (ISAS), as described in Gaillard et al. (2016) is a data based reanalysis of temperature and salinity fields over the global ocean. It was initially designed to synthesize the temperature and salinity profiles collected by the Argo program. It has been later extended to accommodate all type of vertical profile as well as time series. ISAS gridded fields are entirely based on in-situ measurements. The methodology and configuration have been conceived to preserve as much as possible the data information content and resolution. ISAS is developed and run in a research laboratory (LOPS) in close collaboration with Coriolis, one of Argo Global Data Assembly Center and unique data provider for the Mercator operational oceanography system. At the moment the period covered starts in 2002 and only the upper 2000 m are considered. The gridded fields were produced over the global ocean $70^{\circ}N-70^{\circ}S$ on a $1/2^{\circ}$ grid by the ISAS project with datasets downloaded from the Coriolis data center (for more details on ISAS see Gaillard et al. (2009)). In the Pi-MEP, the product in used is the INSITU_GLO_TS_OA_NRT_OBSERVATIONS_013_002_a v6.2 NRT derived at the Coriolis data center and provided by Copernicus (www.marine.copernicus.eu/documents/ PUM/CMEMS-INS-PUM-013-002-ab.pdf). The major contribution to the data set is from Argo array of profiling floats, reaching an approximate resolution of one profile every 10-days and every 3-degrees over the satellite SSS period (http://www.umr-lops.fr/SNO-Argo/Products/ ISAS-T-S-fields/); in this version SSS from ship of opportunity thermosalinographs are not used, so that we can consider SMOS SSS validation using these measurements independent of ISAS. The ISAS optimal interpolation involves a structure function modeled as the sum of two Gaussian functions, each associated with specific time and space scales, resulting in a smoothing over typically 3 degrees. The smallest scale which can be retrieved with ISAS analysis is not smaller than 300–500 km (Kolodziejczyk et al. (2015)). For validation purpose, the ISAS monthly SSS fields at 5 m depth are collocated and compared with the satellite SSS products and included in the Pi-MEP Match-up files. In addition, the "percentage of variance" fields (PCTVAR) contained in the ISAS analyses provide information on the local variability of in situ SSS measurements within $1/2^{\circ} \times 1/2^{\circ}$ boxes.



2.3.4 World Ocean Atlas Climatology

The World Ocean Atlas 2013 version 2 (WOA13 V2) is a set of objectively analyzed (1° grid) climatological fields of in situ temperature, salinity and other variables provided at standard depth levels for annual, seasonal, and monthly compositing periods for the World Ocean. It also includes associated statistical fields of observed oceanographic profile data interpolated to standard depth levels on 5°, 1°, and 0.25° grids. We use these fields in complement to ISAS to characterize the climatological fields (annual mean and std) at the match-up pairs location and date.

2.4 Overview of the Match-ups generation method

The match-up production is basically a three steps process:

- 1. preparation of the input in situ and satellite data, and,
- 2. co-localization of satellite products with in situ SSS measurements.
- 3. co-localization of the in situ/satellite pair with auxiliary information.

In the following, we successively detail the approaches taken for these different steps.

2.4.1 In Situ/Satellite data filtering

The first step consist in filtering TSG-LEGOS-DMin situ dataset using the quality flags as described in 2.2 so that only valid salinity data remains in the produced match-ups.

For high-spatial resolution in situ SSS measurements such as the Thermo-SalinoGraph (TSG) SSS data from research vessels, Voluntary Observing Ships (VOS) or sailing ships, as well as SSS data from surface drifters, an additional spatial-filtering step is performed on the in situ data that will be in fine compared to the satellite SSS products. If R_{sat} is the spatial resolution of the satellite SSS product (L2 to L3-L4), we keep the in situ data at the original spatial resolution but we also estimate for all spatio-temporal samples a running median filtered SSS applied to all neighbouring in situ SSS data acquired within a distance of $R_{sat}/2$ from a given in situ acquisition. Both the original and the filtered data are finally stored in the MDB files.

Only for satellite L2 SSS data, a third step consist in filtering spurious data using the flags and associated recommendation as provided by the official data centers and described in 2.1.

2.4.2 In Situ/Satellite Co-localization

In this step, each SSS satellite acquisition is co-localized with the filtered in situ measurements. The method used for co-localization differ if the satellite SSS is a swath product (so-called Level 2-types) or a time-space composite product (so-called Level 3/level 4-types).

• For L2 SSS swath data:

If R_{sat} is the spatial resolution of the satellite swath SSS product, for each in situ data sample collected in the Pi-MEP database, the platform searches for all satellite SSS data found at grid nodes located within a radius of $R_{sat}/2$ from the in situ data location and acquired with a time-lag from the in situ measurement date that is less or equal than \pm 12 hours. If several satellite SSS samples are found to meet these criteria, the final satellite SSS match-up point is selected to be the closest in time from the in situ data measurement date. The final spatial and temporal lags between the in situ and satellite data are stored in the MDB files.



• For L3 and L4 composite SSS products :

If R_{sat} is the spatial resolution of the composite satellite SSS product and D the period over which the composite product was built (e.g., periods of 1, 7, 8, 9, 10, 18 days, 1 month, etc..) with central time to, for each in situ data sample collected in the Pi-MEP database during period D, the platform searches for all satellite SSS data of the composite product found at grid nodes located within a radius of $R_{sat}/2$ from the in situ data location. If several satellite SSS product samples are found to meet these criteria, the final satellite SSS match-up point is chosen to be the composite SSS with central time to which is the closest in time from the in situ data measurement date. The final spatial and temporal lags between the in situ and satellite data are stored in the MDB files.

2.4.3 MDB pair Co-localization with auxiliary data and complementary information

MDB data consist of satellite and in-situ SSS pair datasets but also of auxiliary geophysical parameters such as local and history of wind speed and rain rates, as well as various information (climatology, distance to coast, mixed layer depth, barrier layer thickness, etc) that can be derived from in situ data and which are included in the final match-up files. The collocation of auxiliary parameters and additional information is done for each filtered in-situ SSS measurement contained in the match-up files as follows:

If t_{insitu} is the time/date at which the in situ measurement is performed, we collect:

- The ASCAT wind speed product of the same day than t_{insitu} found at the ASCAT $1/4^{\circ}$ grid node with closest distance from the in situ data location and the time series of the ASCAT wind speed at the same node for the 10 days prior the in situ measurement day.
- If the in situ data is located within the 60°N-60°S band, we select the CMORPH 3-hourly product the closest in time from tin situ and found at the CMORPH 1/4° grid node with closest distance from the in situ data location. We then store the time series of the CMORPH rain rate at the same node for the 10 days prior the in situ measurement time.

For the given month/year of the in situ data, we select the ISAS and WOA fields for the same month (and same year for ISAS fields) and take the SSS analysis (monthly mean, std) found at the closest grid node from the in situ measurement.

The distance from the in situ SSS data location to the nearest coast is evaluated and provided in km. We use a distance-to-coast map at $1/4^{\circ}$ resolution where small islands have been removed.

When vertical profiles of salinity (S) and temperature (T) are made available from the in situ measurements used to build the match-up (Argo or sea mammals), the following variables are included into each satellite/in situ match-up file:

- 1. The vertical distribution of pressure at which the profile were measured,
- 2. The vertical S(z) and T(z) profiles,
- 3. The vertical potential density anomaly profile $\sigma_0(z)$,
- 4. The Mixed Layer Depth (MLD). The MLD is defined here as the depth where the potential density has increased from the reference depth (10 meter) by a threshold equivalent to 0.2° C decrease in temperature at constant salinity: $\sigma_0 = \sigma_{010m} + \Delta \sigma_0$ with $\Delta \sigma_0 = \sigma_0(\theta_{10m} 0.2, S_{10m}) \sigma_0(\theta_{10m}, S_{10m})$ where θ_{10m} and S_{10m} are the temperature and salinity at the reference depth (i.e. 10 m) (de Boyer Montégut et al. (2004), de Boyer Montégut et al. (2007)).



- 5. The Top of the Thermocline Depth (TTD) is defined as the depth at which temperature decreases from its 10 m value by 0.2° C.
- 6. The Barrier Layer if present, is defined as the intermediate layer between the top of the thermocline and the bottom of the density mixed-layer and its thickness (BLT) is defined as the difference between the MLD and the TTD.
- 7. The vertical profile of the buoyancy frequency $N^2(z)$

The resulting match-ups files are serialized as NetCDF-4 files whose structure depends on the origin of the in-situ data they contain.

2.4.4 Content of the Match-Up NetCDF files

```
netcdf pimep-mdb_smos-l3-catds-locean-v3-18d_tsg-legos-dm_20100116_v01 {
dimensions:
    TIME\_SAT = UNLIMITED; // (1 currently)
    TIME\_TSG = 2190;
    N_DAYS_WIND = 10;
    N_3H_RAIN = 80;
    STRING25 = 25;
    STRING8 = 8;
   variables:
float DATE_TSG(TIME_TSG) ;
DATE\_TSG:long\_name = "Date of TSG";
    DATE_TSG:units = "days since 1990-01-01 00:00:00";
    DATE_TSG:standard_name = "time";
    DATE_TSG:_FillValue = -999.f;
float LATITUDE_TSG(TIME_TSG);
    LATITUDE_TSG:long_name = "Latitude of TSG";
    LATITUDE\_TSG:units = "degrees\_north" \ ;
    LATITUDE_TSG:valid_min = -90. :
    LATITUDE_TSG:valid_max = 90.;
    LATITUDE\_TSG: standard\_name = "latitude" \ ;
    LATITUDE_TSG:_FillValue = -999.f;
float LONGITUDE_TSG(TIME_TSG);
    LONGITUDE_TSG:long_name = "Longitude of TSG";
    LONGITUDE_TSG:units = "degrees_east";
    LONGITUDE_TSG:valid_min = -180.;
    LONGITUDE\_TSG:valid\_max = 180.;
    LONGITUDE_TSG:standard_name = "longitude";
    LONGITUDE\_TSG:\_FillValue = -999.f;
float SSS_TSG(TIME_TSG);
    SSS_TSG:long_name = "TSG SSS" :
    SSS_TSG:units = "1";
    SSS_TSG:salinity_scale = "Practical Salinity Scale(PSS-78)";
    SSS_TSG:standard_name = "sea_water_salinity";
    SSS\_TSG:\_FillValue = -999.f;
float SST_TSG(TIME_TSG);
```



```
SST_TSG:long_name = "TSG SST";
    SST_TSG:units = "degree Celsius";
    SST_TSG:standard_name = "sea_water_temperature";
    SST_TSG:_FillValue = -999.f;
float SSS_TSG_FILTERED(TIME_TSG) ;
    SSS_TSG_FILTERED:long_name = "TSG SSS median filtered at satellite spatial resolu-
tion";
    SSS\_TSG\_FILTERED:units = "1";
    SSS_TSG_FILTERED:salinity_scale = "Practical Salinity Scale(PSS-78)";
    SSS_TSG_FILTERED:standard_name = "sea_water_salinity";
    SSS\_TSG\_FILTERED:\_FillValue = -999.f;
float SST_TSG_FILTERED(TIME_TSG);
    SST_TSG_FILTERED:long_name = "TSG SST median filtered at satellite spatial resolu-
tion";
    SST_TSG_FILTERED:units = "degree Celsius";
    SST_TSG_FILTERED:standard_name = "sea_water_temperature";
    SST_TSG_FILTERED:_FillValue = -999.f;
float DISTANCE_TO_COAST_TSG(TIME_TSG);
    DISTANCE_TO_COAST_TSG:long_name = "Distance to coasts at TSG location";
    DISTANCE_TO_COAST_TSG:units = "km";
    DISTANCE_TO_COAST_TSG:_FillValue = -999.f;
float PLATFORM_NUMBER_TSG(TIME_TSG);
    PLATFORM_NUMBER_TSG:long_name = "TSG unique identifier";
    PLATFORM_NUMBER_TSG:conventions = "WMO float identifier : A9IIIII";
    PLATFORM_NUMBER_TSG:units = "1";
    PLATFORM_NUMBER_TSG:_FillValue = -999.f;
float DATE_Satellite_product(TIME_Sat);
    DATE_Satellite_product:long_name = "Central time of satellite SSS file";
    DATE\_Satellite\_product:units = "days since 1990-01-01 00:00:00" ;
    DATE_Satellite_product:standard_name = "time";
float LATITUDE_Satellite_product(TIME_TSG) ;
    LATITUDE_Satellite_product:long_name = "Satellite product latitude at TSG location";
    LATITUDE_Satellite_product:units = "degrees_north";
    LATITUDE_Satellite_product:valid_min = -90. ;
    LATITUDE_Satellite_product:valid_max = 90.;
    LATITUDE\_Satellite\_product:standard\_name = "latitude" \ ;
    LATITUDE_Satellite_product:_FillValue = -999.f;
float LONGITUDE_Satellite_product(TIME_TSG);
    LONGITUDE_Satellite_product:long_name = "Satellite product longitude at TSG location"
    LONGITUDE_Satellite_product:units = "degrees_east";
    LONGITUDE_Satellite_product:valid_min = -180. :
    LONGITUDE_Satellite_product:valid_max = 180.;
    LONGITUDE_Satellite_product:standard_name = "longitude";
    LONGITUDE_Satellite_product:_FillValue = -999.f;
SSS\_Satellite\_product:units = "1";
    SSS_Satellite_product:salinity_scale = "Practical Salinity Scale(PSS-78)";
```



```
SSS_Satellite_product:standard_name = "sea_surface_salinity";
    SSS\_Satellite\_product:\_FillValue = -999.f;
float SST_Satellite_product(TIME_TSG);
    SST_Satellite_product:long_name = "Satellite product SST at TSG location";
    SST_Satellite_product:units = "degree Celsius";
    SST_Satellite_product:standard_name = "sea_surface_temperature";
    SST\_Satellite\_product:\_FillValue = -999.f;
float Spatial_lags(TIME_TSG) ;
    Spatial_lags:long_name = "Spatial lag between TSG location and satellite SSS product pixel
center":
    Spatial_lags:units = "km";
    Spatial_lags:_FillValue = -999.f;
float Time_lags(TIME_TSG);
    Time_lags:long_name = "Temporal lag between TSG time and satellite SSS product central
time":
    Time_{-lags:units} = "days";
    Time_{lags:}FillValue = -999.f;
float ROSSBY_RADIUS_at_TSG(TIME_TSG);
    ROSSBY_RADIUS_at_TSG:long_name = "Baroclinic Rossby radius of deformation (Chel-
ton et al., 1998) at TSG location";
    ROSSBY_RADIUS_at_TSG:units = "km";
    ROSSBY_RADIUS_at_TSG:_FillValue = -999.f;
float Ascat_daily_wind_at_TSG(TIME_TSG);
    Ascat_daily_wind_at_TSG:long_name = "Daily Ascat wind speed module at TSG location"
    Ascat_daily_wind_at_TSG:units = "m/s";
    Ascat_daily\_wind_at_TSG:_FillValue = -999.f;
float CMORPH_3h_Rain_Rate_at_TSG(TIME_TSG) ;
     CMORPH_3h_Rain_Rate_at_TSG:long_name = "3-hourly CMORPH rain rate at TSG lo-
cation";
    CMORPH_3h_Rain_Rate_at_TSG:units = "mm/3h";
    CMORPH_3h_Rain_Rate_at_TSG:_FillValue = -999.f;
float Ascat_10_prior_days_wind_at_TSG(TIME_TSG, N_DAYS_WIND);
     Ascat_10_prior_days_wind_at_TSG:long_name = "Prior 10 days time series of Ascat wind
speed module at TSG location";
    Ascat_10_prior_days_wind_at_TSG:units = "m/s";
    Ascat_10\_prior_days\_wind_at_TSG:_FillValue = -999.f;
float CMORPH_10_prior_days_Rain_Rate_at_TSG(TIME_TSG, N_3H_RAIN);
    CMORPH_10_prior_days_Rain_Rate_at_TSG:long_name = "Prior 10 days times series of 3-
hourly CMORPH Rain Rate at TSG location";
    CMORPH_10_prior_days_Rain_Rate_at_TSG:units = "mm/3h";
    CMORPH_10_prior_days_Rain_Rate_at_TSG:_FillValue = -999.f;
float SSS_ISAS_at_TSG(TIME_TSG) ;
    SSS_ISAS_at_TSG:long_name = "ISAS SSS (5m depth) at TSG location";
    SSS_ISAS_at_TSG:units = "1";
    SSS_ISAS_at_TSG:salinity_scale = "Practical Salinity Scale(PSS-78)";
    SSS_ISAS_at_TSG:standard_name = "sea_water_salinity";
    SSS\_ISAS\_at\_TSG:\_FillValue = -999.f;
float SSS_PCTVAR_ISAS_at_TSG(TIME_TSG);
```



```
SSS_PCTVAR_ISAS_at_TSG:long_name = "Error on ISAS SSS (5m depth) at TSG location
(% variance)";
    SSS_PCTVAR_ISAS_at_TSG:units = "%";
    SSS_PCTVAR_ISAS_at_TSG:_FillValue = -999.f;
float SSS_WOA13_at_TSG(TIME_TSG);
    SSS_WOA13_at_TSG:long_name = "WOA 2013 (DECAV-1deg) SSS (0m depth) at TSG
location"
    SSS_WOA13_at_TSG:units = "1";
    SSS_WOA13_at_TSG:salinity_scale = "Practical Salinity Scale(PSS-78)";
    SSS_WOA13_at_TSG:standard_name = "sea_surface_salinity";
    SSS_WOA13_at_TSG:_FillValue = -999.f;
float SSS_STD_WOA13_at_TSG(TIME_TSG);
    SSS_STD_WOA13_at_TSG:long_name = "WOA 2013 (DECAV-1deg) SSS STD (0m depth)
at TSG location";
    SSS\_STD\_WOA13\_at\_TSG:units = "1";
    SSS\_STD\_WOA13\_at\_TSG:\_FillValue = -999.f;
float SSS_ISAS15_at_TSG(N_prof);
    SSS_ISAS15_at_TSG:long_name = "Monthly ISAS-15 SSS (5m depth) at TSG location";
    SSS\_ISAS15\_at\_TSG:units = "1";
    SSS_ISAS15_at_TSG:salinity_scale = "Practical Salinity Scale (PSS-78)";
    SSS_ISAS15_at_TSG:standard_name = "sea_water_salinity";
    SSS_ISAS15_at_TSG:_FillValue = -999.f;
float SSS_PCTVAR_ISAS15_at_TSG(N_prof);
    SSS_PCTVAR_ISAS15_at_TSG:long_name = "Error on monthly ISAS-15 SSS (5m depth)
at TSG location (% variance)";
    SSS_PCTVAR_ISAS15_at_TSG:units = "%";
    SSS_PCTVAR_ISAS15_at_TSG:_FillValue = -999.f;
float SSS_WOA18_at_TSG(N_prof);
    SSS_WOA18_at_TSG:long_name = "Monthly WOA 2018 (DECAV-1deg) SSS (0m depth)
at TSG location";
    SSS_WOA18_at_TSG:units = "1";
    SSS_WOA18_at_TSG:salinity_scale = "Practical Salinity Scale (PSS-78)";
    SSS_WOA18_at_TSG:standard_name = "sea_surface_salinity";
    SSS_WOA18_at_TSG:_FillValue = -999.f;
float SSS_STD_WOA18_at_TSG(N_prof);
    SSS\_STD\_WOA18\_at\_TSG:long\_name = "Monthly WOA 2018 (DECAV-1deg) SSS STD
(0m depth) at TSG location";
    SSS\_STD\_WOA18\_at\_TSG:units = "1";
    SSS\_STD\_WOA18\_at\_TSG:\_FillValue = -999.f;
float SEA_ICE_CONCENTRATION_at_TSG(N_prof);
    SEA_ICE_CONCENTRATION_at_TSG:long_name = "Daily sea ice area fraction (EUMET-
SAT OSI-SAF OSI-450) at TSG location (%)":
    SEA_ICE_CONCENTRATION_at_TSG:units = "1";
    SEA_ICE_CONCENTRATION_at_TSG:standard_name = "sea_ice_area_fraction";
    SEA_ICE_CONCENTRATION_at_TSG:_FillValue = -999.f;
float CCMP_6h_Wind_Speed_at_TSG(N_prof);
    CCMP_6h_Wind_Speed_at_TSG:long_name = "6-hourly CCMP wind speed at TSG loca-
tion";
    CCMP_6h_Wind_Speed_at_TSG:units = "m s-1";
```



```
CCMP_6h_Wind_Speed_at_TSG:standard_name = "wind_speed";
         CCMP_6h_Wind_Speed_at_TSG:_FillValue = -999.f;
float CCMP_10_prior_days_Wind_Speed_at_TSG(N_prof, N_DAYS_WIND_CCMP);
        CCMP_10_prior_days_Wind_Speed_at_TSG:long_name = "Prior 10 days time series of CCMP
wind speed at TSG location";
         CCMP_10_prior_days_Wind_Speed_at_TSG:units = "m s-1";
         CCMP\_10\_prior\_days\_Wind\_Speed\_at\_TSG:standard\_name = "wind\_speed" \ ;
         CCMP_10_prior_days_Wind_Speed_at_TSG:_FillValue = -999.f;
float CDM_GLOBCOLOUR_at_TSG(N_prof);
         CDM_GLOBCOLOUR_at_TSG:long_name = "8-day Coloured dissolved and detrital or-
ganic materials - mean of the binned pixels at TSG location";
         CDM_GLOBCOLOUR_at_TSG:units = "m-1";
        CDM\_GLOBCOLOUR\_at\_TSG: standard\_name = "volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_flux\_in\_sea\_water\_radiative\_f
         CDM\_GLOBCOLOUR\_at\_TSG:\_FillValue = -999.f;
float CHL1_GLOBCOLOUR_at_TSG(N_prof);
         CHL1_GLOBCOLOUR_at_TSG:long_name = "8-day Chlorophyll concentration - mean of
the binned pixels at TSG location";
         CHL1_GLOBCOLOUR_at_TSG:units = "mg m-3";
        CHL1_GLOBCOLOUR_at_TSG:standard_name = "mass_concentration_of_chlorophyll_a_in_sea_water"
         CHL1_GLOBCOLOUR_at_TSG:_FillValue = -999.f;
float EVAPORATION_OAFLUX_at_TSG(N_prof) ;
        EVAPORATION_OAFLUX_at_TSG:long_name = "Daily mean evaporation rate (OAFlux)
at TSG location";
         EVAPORATION_OAFLUX_at_TSG:units = "cm year-1";
         EVAPORATION_OAFLUX_at_TSG:_FillValue = -999.f;
float SSS_SCRIPPS_at_TSG(N_prof);
         SSS_SCRIPPS_at_TSG:long_name = "Argo gridded monthly mean SSS (0m depth) from
SCRIPPS (Roemmich-Gilson) at TSG location";
         SSS\_SCRIPPS\_at\_TSG:units = "1";
         SSS_SCRIPPS_at_TSG:salinity_scale = "Practical Salinity Scale (PSS-78)";
         SSS_SCRIPPS_at_TSG:standard_name = "sea_water_salinity";
         SSS\_SCRIPPS\_at\_TSG:\_FillValue = -999.f;
float SSS_IPRC_at_TSG(N_prof);
        SSS_IPRC_at_TSG:long_name = "Argo gridded monthly mean SSS (0m depth) from IPRC
at TSG location";
         SSS\_IPRC\_at\_TSG:units = "1";
         SSS_IPRC_at_TSG:salinity_scale = "Practical Salinity Scale (PSS-78)";
        SSS_IPRC_at_TSG:standard_name = "sea_water_salinity";
         SSS\_IPRC\_at\_TSG:\_FillValue = -999.f;
float SST_AVHRR_at_TSG(N_prof);
         SST_AVHRR_at_TSG:long_name = "Daily OI AVHRR-only v2 SST (Reynolds et al., 2007)
at TSG location";
         SST_AVHRR_at_TSG:units = "degree Celsius";
         SST_AVHRR_at_TSG:standard_name = "sea_water_temperature";
         SST_AVHRR_at_TSG:_FillValue = -999.f;
float \ U\_EKMAN\_GLOBCURRENT\_at\_TSG(N\_prof) \ ;
         U_EKMAN_GLOBCURRENT_at_TSG:long_name = "15m depth Ekman current velocity:
```



```
zonal component at TSG location";
    U_EKMAN_GLOBCURRENT_at_TSG:units = "m s-1";
    U_EKMAN_GLOBCURRENT_at_TSG:_FillValue = -999.f;
float V_EKMAN_GLOBCURRENT_at_TSG(N_prof);
    V_EKMAN_GLOBCURRENT_at_TSG:long_name = "15m depth Ekman current velocity:
meridian component at TSG location";
    V_EKMAN_GLOBCURRENT_at_TSG:units = "m s-1";
    V_EKMAN_GLOBCURRENT_at_TSG:_FillValue = -999.f;
float U_GEOSTROPHIC_GLOBCURRENT_at_TSG(N_prof);
    U_GEOSTROPHIC_GLOBCURRENT_at_TSG:long_name = "Absolute geostrophic veloc-
ity: zonal component at TSG location";
    U_GEOSTROPHIC_GLOBCURRENT_at_TSG:units = "m s-1";
    U_GEOSTROPHIC_GLOBCURRENT_at_TSG:_FillValue = -999.f;
float V_GEOSTROPHIC_GLOBCURRENT_at_TSG(N_prof);
    V_GEOSTROPHIC_GLOBCURRENT_at_TSG:long_name = "Absolute geostrophic veloc-
ity: meridian component at TSG location";
    V_GEOSTROPHIC_GLOBCURRENT_at_TSG:units = "m s-1";
    V_GEOSTROPHIC_GLOBCURRENT_at_TSG:_FillValue = -999.f;
   // global attributes:
    :Conventions = "CF-1.6";
    :title = "TSG-LEGOS-DM Match-Up Database";
    : Satellite\_product\_name = "SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM" \ ;
    :Satellite_product_spatial_resolution = "25 km";
    :Satellite_product_temporal_resolution = "18 days";
    :Satellite_product_filename = "v3/18days/SMOS_L3_DEBIAS_LOCEAN_AD_20100116_EASE_18d_25km_v03.nc";
    :Match-Up_spatial_window_radius_in_km = 25;
    :Match-Up\_temporal\_window\_radius\_in\_days = 2;
    :start\_time = "20100114T000005Z";
    :stop\_time = "20100118T235026Z";
    :northernmost_latitude = 77.676f;
    :sourthenmost_latitude = -66.423f;
    :westernmost_longitude = -179.219f;
    :easternmost_longitude = 179.199f;
    :geospatial_lat_units = "degrees north";
    :geospatial_lat_resolution = "25 km";
    :geospatial_lon_units = "degrees east";
    :geospatial_lon_resolution = "25 km"
    : institution = "ESA-IFREMER-ODL" ; \\
    :project_name = "SMOS Pilote Mission Exploitation Platfrom (Pi-MEP) for salinity";
    :project_url = "https://pimep-project.odl.bzh";
    :license = "Pi-MEP data use is free and open" :
    :product_version = "1.0";
    :keywords = "Oceans > Ocean Salinity > Sea Surface Salinity";
    :acknowledgment = "Please acknowledge the use of these data with the following statement:
These data were provided by SMOS Pilote Mission Exploitation Platfrom (Pi-MEP) for salinity"
    :Source = "v3/18days/SMOS_L3_DEBIAS_LOCEAN_AD_20100116_EASE_18d_25km_v03.nc";
    :references = "https://pimep-project.odl.bzh";
```



```
:history = "Processed on 2018-04-18 using MDB_generator" ; :date_created = "2018-04-18 17:09:30" ; }
```

2.5 MDB characteristics for the particular in situ/satellite pairs

2.5.1 Number of paired SSS data as a function of time and distance to coast




Figure 1: Number of match-ups between TSG-LEGOS-DM and SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM SSS as a function of time (a) and as function of the distance to coast (b) over the North Atlantic Ocean Pi-MEP region and for the full satellite product period.

2.5.2 Histograms of the SSS match-ups




Figure 2: Histograms of SSS from TSG-LEGOS-DM (a) and SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM (b) considering all match-up pairs per bins of 0.1 over the North Atlantic Ocean Pi-MEP region and for the full satellite product period.



2.5.3 Distribution of in situ SSS depth measurements




Figure 3: Histograms of the depth of the upper level SSS measurements from TSG-LEGOS-DM in the Match-up DataBase for the North Atlantic Ocean Pi-MEP region (a) and temporal mean spatial distribution of pressure of the in situ SSS data over 1°x1° boxes and for the full satellite product period (b).

2.5.4 Spatial Distribution of Match-ups




Figure 4: Number of SSS match-ups between TSG-LEGOS-DM SSS and the SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM SSS product for the North Atlantic Ocean Pi-MEP region over $1^{\circ}x1^{\circ}$ boxes and for the full satellite product period.



2.5.5 Histograms of the spatial and temporal lags of the match-ups pairs




Figure 5: Histograms of the spatial (a) and temporal (b) lags between the time of the TSG-LEGOS-DM measurements and the date of the corresponding SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM SSS product.

3 MDB file Analyses

3.1 Spatial Maps of the Temporal mean and Std of in situ and satellite SSS and of the difference (Δ SSS)

In Figure 6, we show maps of temporal mean (left) and standard deviation (right) of the SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM satellite SSS product (top) and of the TSG-LEGOS-DM in situ dataset at the collected Pi-MEP match-up pairs. The temporal mean and std are gridded over the full satellite product period and over spatial boxes of size 1°x1°.

At the bottom of Figure 6, the temporal mean (left) and standard deviation (right) of the differences between the satellite SSS product and in situ data found at match-up pairs, namely Δ SSS(Satellite -TSG-LEGOS-DM), is also gridded over the full satellite product period and over spatial boxes of size 1°x1°.






Figure 6: Temporal mean (left) and Std (right) of SSS from SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM (top), TSG-LEGOS-DM (middle), and of Δ SSS (Satellite - TSG-LEGOS-DM). Only match-up pairs are used to generate these maps.



3.2 Time series of the monthly averaged mean and Std of in situ and satellite SSS and of the (Δ SSS)

In the top panel of Figure 7, we show the time series of the monthly averaged SSS estimated over the full North Atlantic Ocean Pi-MEP region for both SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM satellite SSS product (in black) and the TSG-LEGOS-DM in situ dataset (in blue) at the collected Pi-MEP match-up pairs.

In the middle panel of Figure 7, we show the time series of the monthly averaged Δ SSS (Satellite - TSG-LEGOS-DM) for the collected Pi-MEP match-up pairs and estimated over the full North Atlantic Ocean Pi-MEP region.

In the bottom panel of Figure 7, we show the time series of the monthly averaged standard deviation of the Δ SSS (Satellite - TSG-LEGOS-DM) for the collected Pi-MEP match-up pairs and estimated over the full North Atlantic Ocean Pi-MEP region.




Figure 7: Time series of the monthly averaged mean SSS (top), mean Δ SSS (Satellite - TSG-LEGOS-DM) and Std of Δ SSS (Satellite - TSG-LEGOS-DM) over the North Atlantic Ocean Pi-MEP region considering all match-ups collected by the Pi-MEP platform.

3.3 Zonally-averaged Time-mean and temporal Std of in situ and satellite SSS and of the Δ SSS

In Figure 8 left panel, we show the zonally averaged time-mean SSS estimated at the collected Pi-MEP match-up pairs for both SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM satellite SSS product (in black) and the TSG-LEGOS-DM in situ dataset (in blue). The time mean is evaluated over the full satellite SSS product period.

In the right panel of Figure 8, we show the zonally averaged time-mean Δ SSS (Satellite - TSG-LEGOS-DM) for all the collected Pi-MEP match-up pairs estimated over the full satellite product period.






Figure 8: Left panel: Zonally averaged time mean SSS from SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM (black) and from TSG-LEGOS-DM (blue). Right panel: zonally averaged time-mean Δ SSS (Satellite - TSG-LEGOS-DM) for all the collected Pi-MEP match-up pairs estimated over the full satellite product period.



3.4 Scatterplots of satellite vs in situ SSS by latitudinal bands




Figure 9: Contour maps of the concentration of SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM SSS (y-axis) versus TSG-LEGOS-DM SSS (x-axis) at match-up pairs for different latitude bands. For each plot, the red line shows x=y. The black thin and dashed lines indicate a linear fit through the data cloud and the $\pm 95\%$ confidence levels, respectively. The number match-up pairs n, the slope and \mathbf{R}^2 coefficient of the linear fit, the root mean square (RMS) and the mean bias between satellite and in situ data are indicated for each latitude band in each plots.



3.5 Time series of the monthly averaged mean and Std of the ΔSSS sorted by latitudinal bands




Figure 10: Monthly-average mean (red curves) Δ SSS (Satellite - TSG-LEGOS-DM) and ± 1 Std (black vertical thick bars) as function of time for all the collected Pi-MEP match-up pairs estimated over the North Atlantic Ocean Pi-MEP region and for the full satellite product period are shown for different latitude bands: (a) Latitude band 80°S-80°N, (b) latitude band 20°S-20°N, (c) Mid Latitude bands 40°S-20°S and 20°N-40°N and (d) Latitude bands 60°S-40°S and 40°N-60°N.

3.6 Δ SSS sorted as function of geophysical parameters

In Figure 11, we classify the match-up differences Δ SSS (Satellite - in situ) between SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM and TSG-LEGOS-DM SSS as function of the geophysical conditions at match-up points. The mean and std of Δ SSS (Satellite - TSG-LEGOS-DM) is thus evaluated as function of the

- in situ SSS values per bins of width 0.2,
- in situ SST values per bins of width 1°C,
- ASCAT daily wind values per bins of width 1 m/s,
- CMORPH 3-hourly rain rates per bins of width 1 mm/h, and,
- distance to coasts per bins of width 50 km.






Figure 11: Δ SSS (Satellite - TSG-LEGOS-DM) sorted as function of TSG-LEGOS-DM SSS values a), TSG-LEGOS-DM SST b), ASCAT Wind speed c), CMORPH rain rate d) and distance to coast (e). In all plots the mean and Std of Δ SSS for each bin is indicated by the red curves and black vertical thick bars (± 1 Std)

3.7 Δ SSS maps and statistics for different geophysical conditions

In Figures 12 and 13, we focus on sub-datasets of the match-up differences Δ SSS (Satellite - in situ) between SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM and TSG-LEGOS-DM for the following specific geophysical conditions:

- C1:if the local value at in situ location of estimated rain rate is zero, mean daily wind is in the range [3, 12] m/s, the SST is > 5°C and distance to coast is > 800 km.
- C2:if the local value at in situ location of estimated rain rate is zero, mean daily wind is



in the range [3, 12] m/s.

- C3:if the local value at in situ location of estimated rain rate is high (ie. > 1 mm/h) and mean daily wind is low (ie. < 4 m/s).
- C5:if the in situ data is located where the climatological SSS standard deviation is low (ie. above < 0.2).
- C6:if the in situ data is located where the climatological SSS standard deviation is high (ie. above > 0.2).

For each of these conditions, the temporal mean (gridded over spatial boxes of size $1^{\circ}x1^{\circ}$) and the histogram of the difference ΔSSS (Satellite - in situ) are presented.




Figure 12: Temporal mean gridded over spatial boxes of size 1°x1° of Δ SSS (SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM - TSG-LEGOS-DM) for 5 different subdatasets corresponding to:RR=0 mm/h, 3< U_{10} <12 m/s, SST>5°C, distance to coast > 800 km (a), RR=0 mm/h, 3< U_{10} <12 m/s (b), RR>1mm/h and U_{10} <4m/s (c), WOA2013 SSS Std<0.2 (d), WOA2013 SSS Std>0.2 (e).






Figure 13: Normalized histogram of Δ SSS (SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM - TSG-LEGOS-DM) for 5 different subdatasets corresponding to: RR=0 mm/h, 3< U_{10} <12 m/s, SST>5°C, distance to coast > 800 km (a), RR=0 mm/h, 3< U_{10} <12 m/s (b), RR>1mm/h and U_{10} <4m/s (c), WOA2013 SSS Std<0.2 (d), WOA2013 SSS Std>0.2 (e).

4 Summary

Table 1 shows the mean, median, standard deviation (Std), root mean square (RMS), interquartile range (IQR), correlation coefficient ($\rm r^2$) and robust standard deviation (Std*) of the match-up differences Δ SSS (Satellite - in situ) between SMOS-L3-CATDS-CECOS-LOCEAN-V3-18DAYS-25KM and TSG-LEGOS-DM derived over the North Atlantic Ocean Pi-MEP region and for the full satellite product period and for the following conditions:

- all: All the match-up pairs satellite/in situ SSS values are used to derive the statistics
- C1: only pairs where RR=0 mm/h, $3 < U_{10} < 12$ m/s, SST>5°C, distance to coast > 800 km
- \bullet C2: only pairs where RR=0 mm/h, 3 < U_{10} <12 m/s
- C3: only pairs where RR>1mm/h and U_{10} <4m/s
- C5: only pairs where WOA2013 SSS Std<0.2
- C6: only pairs at WOA2013 SSS Std>0.2
- C7a: only pairs with a distance to coast < 150 km.
- C7b: only pairs with a distance to coast in the range [150, 800] km.
- C7c: only pairs with a distance to coast > 800 km.
- C8a: only pairs where SST is $< 5^{\circ}$ C.
- C8b: only pairs where SST is in the range [5, 15]°C.



- C8c: only pairs where SST is > 15°C.
- C9a: only pairs where SSS is < 33.
- C9b: only pairs where SSS is in the range [33, 37].
- C9c: only pairs where SSS is > 37.

Table 1: Statistics of Δ SSS (Satellite - TSG-LEGOS-DM)

Condition	#	Median	Mean	\mathbf{Std}	RMS	IQR	\mathbf{r}^2	\mathbf{Std}^{\star}
all	2198356	-0.04	0.07	0.91	0.91	0.52	0.70	0.39
C1	168171	-0.02	-0.02	0.20	0.20	0.23	0.93	0.17
C2	1188947	-0.05	0.03	0.89	0.89	0.44	0.72	0.33
C3	7736	-0.04	0.18	1.16	1.18	0.58	0.79	0.42
C5	1105893	-0.07	-0.11	0.38	0.40	0.42	0.80	0.31
C6	940336	-0.02	0.21	1.11	1.12	0.70	0.70	0.50
C7a	999456	-0.01	0.26	1.23	1.25	0.89	0.60	0.61
C7b	982394	-0.08	-0.11	0.47	0.48	0.46	0.72	0.34
C7c	216506	-0.02	-0.03	0.22	0.22	0.24	0.94	0.18
C8a	292237	0.27	0.61	1.30	1.43	1.65	0.19	1.09
C8b	1109534	-0.10	-0.04	0.81	0.82	0.58	0.52	0.43
C8c	796585	-0.03	0.02	0.77	0.77	0.34	0.80	0.26
C9a	241781	1.45	1.58	1.69	2.31	1.87	0.22	1.40
C9b	1851204	-0.09	-0.13	0.51	0.53	0.50	0.69	0.37
C9c	105371	-0.04	-0.04	0.16	0.17	0.20	0.55	0.15

For the same conditions, Table 2 presents statistics of Δ SSS (Satellite - ISAS). Only ISAS SSS values with PCTVAR<80% are used to derive the statistics.

Table 2: Statistics of Δ SSS (Satellite - ISAS)

Condition	#	Median	Mean	\mathbf{Std}	RMS	IQR	\mathbf{r}^2	\mathbf{Std}^{\star}
all	1740418	-0.04	-0.08	0.50	0.51	0.43	0.80	0.32
C1	164572	0.00	0.00	0.22	0.22	0.22	0.92	0.16
C2	937863	-0.03	-0.06	0.50	0.50	0.37	0.82	0.28
C3	6060	-0.13	-0.37	1.10	1.16	0.50	0.63	0.38
C5	1020148	-0.06	-0.09	0.36	0.37	0.41	0.82	0.30
C6	641682	-0.03	-0.07	0.62	0.62	0.45	0.81	0.34
C7a	602344	-0.04	-0.05	0.61	0.61	0.52	0.76	0.39
C7b	925889	-0.06	-0.11	0.47	0.48	0.45	0.70	0.33
C7c	212185	-0.01	-0.01	0.23	0.23	0.23	0.93	0.17
C8a	212838	-0.11	-0.09	0.53	0.54	0.61	0.55	0.45
C8b	877913	-0.08	-0.09	0.44	0.45	0.52	0.60	0.39
C8c	649667	-0.01	-0.05	0.57	0.57	0.29	0.77	0.22
C9a	125729	0.08	-0.08	1.20	1.20	0.86	0.35	0.64
C9b	1511398	-0.06	-0.09	0.41	0.42	0.44	0.75	0.32
C9c	103291	0.02	0.02	0.15	0.15	0.20	0.62	0.15

Numerical values can be downloaded as csv files for Table 1 and Table 2.



References

- Gaël Alory, T. Delcroix, P. Téchiné, D. Diverrès, D. Varillon, S. Cravatte, Y. Gouriou, J. Grelet, S. Jacquin, E. Kestenare, and et al. The French contribution to the voluntary observing ships network of sea surface salinity. *Deep-Sea Res. Pt. I*, 105:1–18, November 2015. ISSN 0967-0637. doi: 10.1016/j.dsr.2015.08.005.
- Abderrahim Bentamy and Denis Croize Fillon. Gridded surface wind fields from Metop/ASCAT measurements. *Int. J. Remote Sens.*, 33(6):1729–1754, March 2012. ISSN 1366-5901. doi: 10.1080/01431161.2011.600348.
- Abderrahim Bentamy, Semyon A. Grodsky, James A. Carton, Denis Croizé-Fillon, and Bertrand Chapron. Matching ASCAT and QuikSCAT winds. *J. Geophys. Res.*, 117(C2), February 2012. ISSN 0148-0227. doi: 10.1029/2011JC007479. C02011.
- J. Boutin, J.L. Vergely, S. Marchand, F. D'Amico, A. Hasson, N. Kolodziejczyk, N. Reul, G. Reverdin, and J. Vialard. New SMOS sea surface salinity with reduced systematic errors and improved variability. *Remote Sens. Environ.*, 214:115–134, sep 2018. doi: 10.1016/j.rse.2018.05.022.
- Jacqueline Boutin, Jean-Luc Vergely, and Stéphane Marchand. SMOS SSS L3 debias v2 maps generated by CATDS CEC LOCEAN. V2.1., 2017. doi: 10.17882/52804.
- Jaqueline Boutin, Y. Chao, W. E. Asher, T. Delcroix, R. Drucker, K. Drushka, N. Kolodziejczyk, T. Lee, N. Reul, G. Reverdin, J. Schanze, A. Soloviev, L. Yu, J. Anderson, L. Brucker, E. Dinnat, A. S. Garcia, W. L. Jones, C. Maes, T. Meissner, W. Tang, N. Vinogradova, and B. Ward. Satellite and In Situ Salinity: Understanding Near-Surface Stratification and Sub-footprint Variability. Bull. Am. Meterol. Soc., 97(8):1391–1407, 2016. ISSN 1520-0477. doi: 10.1175/bams-d-15-00032.1.
- Clément de Boyer Montégut, Gurvan Madec, A. S. Fischer, A. Lazar, and D. Ludicone. Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology. *J. Geophys. Res.*, 109(C12):C12003, December 2004. ISSN 0148-0227. doi: 10.1029/2004jc002378.
- Clément de Boyer Montégut, Juliette Mignot, Alban Lazar, and Sophie Cravatte. Control of salinity on the mixed layer depth in the world ocean: 1. General description. *J. Geophys. Res.*, 112(C6):C06011, June 2007. ISSN 0148-0227. doi: 10.1029/2006jc003953.
- Ralph R. Ferraro. SSM/I derived global rainfall estimates for climatological applications. *J. Geophys. Res.*, 1021:16715–16736, 07 1997. doi: 10.1029/97JD01210.
- Ralph R. Ferraro, Fuzhong Weng, Norman C. Grody, and Limin Zhao. Precipitation characteristics over land from the NOAA-15 AMSU sensor. *Geophys. Res. Lett.*, 27(17):2669–2672, 2000. doi: 10.1029/2000GL011665.
- Fabienne Gaillard, E. Autret, V. Thierry, P. Galaup, C. Coatanoan, and T. Loubrieu. Quality Control of Large Argo Datasets. *J. Atmos. Oceanic Technol.*, 26(2):337–351, 2012/10/10 2009. doi: 10.1175/2008JTECHO552.1.
- Fabienne Gaillard, Thierry Reynaud, Virginie Thierry, Nicolas Kolodziejczyk, and Karina von Schuckmann. In Situ-Based Reanalysis of the Global Ocean Temperature and Salinity with ISAS: Variability of the Heat Content and Steric Height. *J. Clim.*, 29(4):1305–1323, February 2016. ISSN 1520-0442. doi: 10.1175/jcli-d-15-0028.1.



- Robert J. Joyce, John E. Janowiak, Phillip A. Arkin, and Pingping Xie. CMORPH: A Method that Produces Global Precipitation Estimates from Passive Microwave and Infrared Data at High Spatial and Temporal Resolution. *J. Hydrometeorol.*, 5(3):487–503, June 2004. ISSN 1525-7541. doi: 10.1175/1525-7541(2004)005(0487:camtpg)2.0.co;2.
- Nicolas Kolodziejczyk, Gilles Reverdin, and Alban Lazar. Interannual Variability of the Mixed Layer Winter Convection and Spice Injection in the Eastern Subtropical North Atlantic. *J. Phys. Oceanogr.*, 45(2):504–525, Feb 2015. ISSN 1520-0485. doi: 10.1175/jpo-d-14-0042.1.
- Christian Kummerow, Y. Hong, W. S. Olson, S. Yang, R. F. Adler, J. McCollum, R. Ferraro, G. Petty, D-B. Shin, and T. T. Wilheit. The Evolution of the Goddard Profiling Algorithm (GPROF) for Rainfall Estimation from Passive Microwave Sensors. *J. Appl. Meteorol.*, 40(11): 1801–1820, 2001. doi: 10.1175/1520-0450(2001)040(1801:TEOTGP)2.0.CO;2.