







Match-up database Analyses Report

Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG

Surface drifters

Tropical Pacific Ocean

prepared by the Pi-MEP Consortium

March 15, 2019

Contents

1	Ove	Overview					
2	The	MDB file datasets	6				
	2.1	Satellite SSS product	6				
		2.1.1 Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG	6				
	2.2	In situ SSS dataset	6				
	2.3	Auxiliary geophysical datasets	7				
		2.3.1 CMORPH	7				
		2.3.2 ASCAT	8				
		2.3.3 ISAS	8				
		2.3.4 World Ocean Atlas Climatology	9				
	2.4	Overview of the Match-ups generation method	9				
		2.4.1 In Situ/Satellite data filtering	9				
		2.4.2 In Situ/Satellite Co-localization	9				
		2.4.3 MDB pair Co-localization with auxiliary data and complementary information	10				
			$10 \\ 11$				
	2.5	*	$17 \\ 17$				
	2.0		17				
		•	17				
			$\frac{1}{18}$				
			18				
		2.0.4 This tograms of the spatial and temporal tags of the matter ups pairs	10				
3	MD	B file Analyses	18				
	3.1	Spatial Maps of the Temporal mean and Std of in situ and satellite SSS and of	10				
	2.0		18				
	3.2	Time series of the monthly averaged mean and Std of in situ and satellite SSS and of the (ASSS)	19				
of the (ΔSSS)							
3.3 Zonally-averaged Time-mean and temporal Std of in situ and satellite SSS the Δ SSS							
	3.4		$\frac{20}{22}$				
	$\frac{3.4}{3.5}$	Time series of the monthly averaged mean and Std of the Δ SSS sorted by latitu-					
	0.0		23				
	3.6		$\frac{23}{23}$				
	$\frac{3.7}{3.7}$		$\frac{20}{24}$				
	0.1	Approximate and statistics for different geophysical conditions					
4	Sun	nmary	2 6				
\mathbf{L}	ist (of Figures					
	1	Number of match-ups between Surface drifters and Aquarius-L3-OR-V5-7DAY-					
		RUNNING-RAIN-MASK-1DEG SSS as a function of time (a) and as function of					
		the distance to coast (b) over the Tropical Pacific Ocean Pi-MEP region and for					
		1 1	17				
	2	Histograms of SSS from Surface drifters (a) and Aquarius-L3-OR-V5-7DAY-RUNNING	} -				
		RAIN-MASK-1DEG (b) considering all match-up pairs per bins of 0.1 over the					
		Tropical Pacific Ocean Pi-MEP region and for the full satellite product period	17				

3	Number of SSS match-ups between Surface drifters SSS and the Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG SSS product for the Tropical Pacific	
4	Ocean Pi-MEP region over 1°x1° boxes and for the full satellite product period. Histograms of the spatial (a) and temporal (b) lags between the time of the Surface drifters measurements and the date of the corresponding Aquarius-L3-OR-V5-	18
5	7DAY-RUNNING-RAIN-MASK-1DEG SSS product	18
	RUNNING-RAIN-MASK-1DEG (top), Surface drifters (middle), and of Δ SSS (Satellite - Surface drifters). Only match-up pairs are used to generate these maps.	19
6	Time series of the monthly averaged mean SSS (top), mean Δ SSS (Satellite - Surface drifters) and Std of Δ SSS (Satellite - Surface drifters) over the Tropical	
	Pacific Ocean Pi-MEP region considering all match-ups collected by the Pi-MEP platform.	20
7	Left panel: Zonally averaged time mean SSS from Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG (black) and from Surface drifters (blue). Right panel: zonally averaged time-mean Δ SSS (Satellite - Surface drifters) for all the	
8	collected Pi-MEP match-up pairs estimated over the full satellite product period. Contour maps of the concentration of Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG SSS (y-axis) versus Surface drifters SSS (x-axis) at match-up pairs	21
	for different latitude bands. For each plot, the red line shows x=y. The black thin and dashed lines indicate a linear fit through the data cloud and the $\pm 95\%$	
	confidence levels, respectively. The number match-up pairs n , the slope and \mathbf{R}^2 coefficient of the linear fit, the root mean square (RMS) and the mean bias between	
9	satellite and in situ data are indicated for each latitude band in each plots Monthly-average mean (red curves) Δ SSS (Satellite - Surface drifters) and ± 1 Std	22
	(black vertical thick bars) as function of time for all the collected Pi-MEP match- up pairs estimated over the Tropical Pacific Ocean Pi-MEP region and for the full satellite product period are shown for different latitude bands: (a) Latitude band 80°S-80°N, (b) latitude band 20°S-20°N, (c) Mid Latitude bands 40°S-20°S and	
	$20^{\circ}\text{N-}40^{\circ}\text{N}$ and (d) Latitude bands $60^{\circ}\text{S-}40^{\circ}\text{S}$ and $40^{\circ}\text{N-}60^{\circ}\text{N}$	23
10	Δ SSS (Satellite - Surface drifters) sorted as function of Surface drifters SSS values a), Surface drifters SST b), ASCAT Wind speed c), CMORPH rain rate d) and distance to coast (e). In all plots the mean and Std of Δ SSS for each bin is	
11	indicated by the red curves and black vertical thick bars (± 1 Std) Temporal mean gridded over spatial boxes of size $1^{\circ}x1^{\circ}$ of Δ SSS (Aquarius-L3-	24
11	OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG - Surface drifters) for 5 different subdatasets corresponding to:RR=0 mm/h, $3 < U_{10} < 12$ m/s, SST>5°C, distance	
	to coast > 800 km (a), RR=0 mm/h, $3 < U_{10} < 12$ m/s (b), RR>1mm/h and	
12	U_{10} <4m/s (c), WOA2013 SSS Std<0.2 (d), WOA2013 SSS Std>0.2 (e) Normalized histogram of Δ SSS (Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASI	25 <mark>K-</mark>
	1DEG - Surface drifters) for 5 different subdatasets corresponding to: RR=0 mm/h, $3 < U_{10} < 12$ m/s, SST>5°C, distance to coast > 800 km (a), RR=0 mm/h, $3 < U_{10} < 12$ m/s (b), RR>1mm/h and $U_{10} < 4$ m/s (c), WOA2013 SSS	
	Std<0.2 (d), WOA2013 SSS Std>0.2 (e)	26



Acronym

Aquarius NASA/CONAE Salinity mission

ASCAT Advanced Scatterometer

ATBD Algorithm Theoretical Baseline Document

BLT Barrier Layer Thickness CMORPH CPC MORPHing technique

CTD Instrument used to measure the conductivity, temperature, and pressure of

seawater

DM Delayed Mode
EO Earth Observation
ESA European Space Agency
FTP File Transfer Protocol

GOSUD Global Ocean Surface Underway Data GTMBA The Global Tropical Moored Buoy Array

Ifremer Institut français de recherche pour l'exploitation de la mer

IPEV Institut polaire français Paul-Émile Victor

IQR Interquartile range ISAS In Situ Analysis System

Kurt Kurtosis (fourth central moment divided by fourth power of the standard de-

viation)

L2 Level 2

LEGOS Laboratoire d'Etudes en Géophysique et Océanographie Spatiales

LOCEAN Laboratoire d'Océanographie et du Climat : Expérimentations et Approches

Numériques

LOPS Laboratoire d'Océanographie Physique et Spatiale

MDB Match-up Data Base

MEOP Marine Mammals Exploring the Oceans Pole to Pole

MLD Mixed Layer Depth

NCEI National Centers for Environmental Information

NRT Near Real Time

NTAS Northwest Tropical Atlantic Station

OI Optimal interpolation

Pi-MEP Pilot Mission Exploitation Platform

PIRATA Prediction and Researched Moored Array in the Atlantic

QC Quality control

 R_{sat} Spatial resolution of the satellite SSS product

RAMA Research Moored Array for African-Asian-Australian Monsoon Analysis and

Prediction

r² Square of the Pearson correlation coefficient

RMS Root mean square

RR Rain rate

SAMOS Shipboard Automated Meteorological and Oceanographic System

Skew Skewness (third central moment divided by the cube of the standard deviation)

SMAP Soil Moisture Active Passive (NASA mission)
SMOS Soil Moisture and Ocean Salinity (ESA mission)
SPURS Salinity Processes in the Upper Ocean Regional Study

SSS Sea Surface Salinity

 SSS_{insitu} In situ SSS data considered for the match-up



 SSS_{SAT} Satellite SSS product considered for the match-up

 Δ SSS Difference between satellite and in situ SSS at colocalized point (Δ SSS =

 SSS_{SAT} - SSS_{insitu})

SST Sea Surface Temperature Std Standard deviation

Std* Robust Standard deviation = median(abs(x-median(x)))/0.67 (less affected by

outliers than Std)

Stratus Surface buoy located in the eastern tropical Pacific

Survostral SURVeillance de l'Océan AuSTRAL (Monitoring the Southern Ocean)

TAO Tropical Atmosphere Ocean

 $TSG \hspace{1cm} ThermoSalinoGraph \\$

WHOI Woods Hole Oceanographic Institution
WHOTS WHOI Hawaii Ocean Time-series Station

WOA World Ocean Atlas



1 Overview

In this report, we present systematic analyses of the Match-up DataBase (MDB) files generated by the Pi-MEP platform within the following Pi-MEP region and for the below pair of Satellite/In situ SSS data:

- Pi-MEP region: Tropical Pacific Ocean (download the corresponding mask here)
- SSS satellite product (SSS_{SAT}): Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG
- In situ dataset (SSS_{Insitu}): Surface drifters (download the corresponding report here)

In the following, $\Delta SSS = SSS_{SAT}$ - SSS_{Insitu} denotes the difference between the satellite and in situ SSS at the colocalized points that form the MDB.

This report presents successively:

The MDB file DataSets (Section 2)

- A short description of the satellite SSS product considered in the match-up (2.1)
- A short description of the In situ SSS dataset considered in the match-up (2.2)
- A short description of the auxiliary geophysical datasets co-localized with SSS pairs (2.3)
- An overview of how the Match-ups were evaluated (2.4)
- An overview of the MDB characteristics for the particular in situ/satellite pairs (2.5)

The major results of the MDB file Analyses (Section 3)

- Spatial Maps of the Time-mean and temporal Std of in situ and satellite SSS and of the Δ SSS (3.1)
- Time series of the monthly averaged mean and Std of in situ and satellite SSS and of the Δ SSS (3.2)
- Zonally-averaged Time-mean and temporal Std of in situ and satellite SSS and of the Δ SSS (3.3)
- Scatterplots of satellite vs in situ SSS by latitudinal bands (3.4)
- Time series of the monthly averaged mean and Std of the Δ SSS sorted by latitudinal bands (3.5)
- Δ SSS sorted as function of geophysical parameters (3.6)
- Δ SSS maps and statistics for different geophysical conditions (3.7)

All analyses are conducted over the Pi-MEP Region specified above and over the full satellite SSS product period.



2 The MDB file datasets

2.1 Satellite SSS product

2.1.1 Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG

Aquarius Level 3 sea surface salinity (SSS) rain-flagged standard mapped image data contains gridded 1 degree spatial resolution SSS averaged over daily, 7 day, monthly, and seasonal time scales. This particular data set is the 7-Day running mean rain-flagged sea surface salinity product for version 5.0 of the Aquarius data set, which is the official end of mission public data release from the AQUARIUS/SAC-D mission. The Aquarius instrument is onboard the AQUARIUS/SAC-D satellite, a collaborative effort between NASA and the Argentinian Space Agency Comision Nacional de Actividades Espaciales (CONAE). The instrument consists of three radiometers in push broom alignment at incidence angles of 29, 38, and 46 degrees incidence angles relative to the shadow side of the orbit. Footprints for the beams are: 76 km (along-track) x 94 km (cross-track), 84 km x 120 km and 96km x 156 km, yielding a total cross-track swath of 370 km. The radiometers measure brightness temperature at 1.413 GHz in their respective horizontal and vertical polarizations (TH and TV). A scatterometer operating at 1.26 GHz measures ocean backscatter in each footprint that is used for surface roughness corrections in the estimation of salinity. The scatterometer has an approximate 390km swath.

Table 1: Satellite SSS product characteristics

Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG						
Spatial resolution	1°x1°					
Temporal repeat	7 Day					
Temporal coverage	From 2011-08-25 to 2015-06-07					
Spatial coverage	Global [-180 180 -90 90]					
Data Provider	NASA Aquarius project, Goddard Space Flight Center, USA					
Release Date	2017-Dec-07					
Version	5					
User Guide	AquariusUserGuide_DatasetV5.0.pdf					
Documentation	ftp://podaac-ftp.jpl.nasa.gov/allData/aquarius/docs/v5/					
DOI	http://dx.doi.org/10.5067/AQR50-3YQCE					

2.2 In situ SSS dataset

The skin depth of the L-band radiometer signal over the ocean is about 1 cm whereas classical surface salinity measured by ships or Argo floats are performed at a few meters depth. In order to improve the knowledge of the SSS variability in the first 50 cm depth, to better document the SSS variability in a satellite pixel and to provide ground-truth as close as possible to the sea surface for validating satellite SSS, the L-band remotely sensed community proposed to deploy numerous surface drifters over various parts of the ocean. Surface drifter data are provided by the LOCEAN (see https://www.locean-ipsl.upmc.fr/smos/drifters/). Only validated data are considered with uncertainty order of 0.01 and 0.1.



2.3 Auxiliary geophysical datasets

Additional EO datasets are used to characterize the geophysical conditions at the in situ/satellite SSS pair measurement locations and time, and 10 days prior the measurements to get an estimate of the geophysical condition and history. As discussed in Boutin et al. (2016), the presence of vertical gradients in, and horizontal variability of, sea surface salinity indeed complicates comparison of satellite and in situ measurements. The additional EO data are used here to get a first estimates of conditions for which L-band satellite SSS measured in the first centimeters of the upper ocean within a 50-150 km diameter footprint might differ from pointwise in situ measurements performed in general between 10 and 5 m depth below the surface. The spatiotemporal variability of SSS within a satellite footprint (50-150 km) is a major issue for satellite SSS validation in the vicinity of river plumes, frontal zones, and significant precipitation. Rainfall can in some cases produce vertical salinity gradients exceeding 1 pss m⁻¹; consequently, it is recommended that satellite and in situ SSS measurements less than 3-6 h after rain events should be considered with care when used in satellite calibration/validation analyses. To identify such situation, the Pi-MEP test platform is first using CMORPH products to characterize the local value and history of rain rate and ASCAT gridded data are used to characterize the local surface wind speed and history. For validation purpose, the ISAS monthly SSS in situ analysed fields at 5 m depth are collocated and compared with the satellite SSS products. The use of ISAS is motivated by the fact that it is used in the SMOS L2 official validation protocol in which systematic comparisons of SMOS L2 retrieved SSS with ISAS are done. In complement to ISAS, monthly std climatological fields from the World Ocean Atlas (WOA13) at the match-up pairs location and date are also used to have an a priori information of the local SSS variability.

2.3.1 CMORPH

Precipitation are estimated using the CMORPH 3-hourly products at 1/4° resolution (Joyce et al. (2004)). CMORPH (CPC MORPHing technique) produces global precipitation analyses at very high spatial and temporal resolution. This technique uses precipitation estimates that have been derived from low orbiter satellite microwave observations exclusively, and whose features are transported via spatial propagation information that is obtained entirely from geostationary satellite IR data. At present NOAA incorporate precipitation estimates derived from the passive microwaves aboard the DMSP 13, 14 and 15 (SSM/I), the NOAA-15, 16, 17 and 18 (AMSU-B), and AMSR-E and TMI aboard NASA's Aqua, TRMM and GPM spacecraft, respectively. These estimates are generated by algorithms of Ferraro (1997) for SSM/I, Ferraro et al. (2000) for AMSU-B and Kummerow et al. (2001) for TMI. Note that this technique is not a precipitation estimation algorithm but a means by which estimates from existing microwave rainfall algorithms can be combined. Therefore, this method is extremely flexible such that any precipitation estimates from any microwave satellite source can be incorporated.

With regard to spatial resolution, although the precipitation estimates are available on a grid with a spacing of 8 km (at the equator), the resolution of the individual satellite-derived estimates is coarser than that - more on the order of 12×15 km or so. The finer "resolution" is obtained via interpolation.

In effect, IR data are used as a means to transport the microwave-derived precipitation features during periods when microwave data are not available at a location. Propagation vector matrices are produced by computing spatial lag correlations on successive images of geostationary satellite IR which are then used to propagate the microwave derived precipitation estimates. This process governs the movement of the precipitation features only. At a given location, the shape and intensity of the precipitation features in the intervening half hour periods between microwave scans are determined by performing a time-weighting interpolation between microwave-derived



features that have been propagated forward in time from the previous microwave observation and those that have been propagated backward in time from the following microwave scan. NOAA refer to this latter step as "morphing" of the features.

For the present Pi-MEP products, we only considered the 3-hourly products at 1/4 degree resolution. The entire CMORPH record (December 2002-present) for 3-hourly, 1/4 degree lat/lon resolution can be found at: ftp://ftp.cpc.ncep.noaa.gov/precip/CMORPH_V1.0/RAW/. CMORPH estimates cover a global belt (-180°W to 180°E) extending from 60°S to 60°N latitude and are available for the complete period of the Pi-MEP core datasets (Jan 2010-now).

2.3.2 ASCAT

Advanced SCATterometer (ASCAT) daily data produced and made available at Ifremer/CERSAT on a 0.25°x0.25° resolution grid (Bentamy and Fillon (2012)) since March 2007 are used to characterize the mean daily wind at the match-up pair location as well as the wind history during the 10-days period preceding the in situ measurement date. These wind fields are calculated based on a geostatistical method with external drift. Remotely sensed data from ASCAT are considered as observations while those from numerical model analysis (ECMWF) are associated with the external drift. The spatial and temporal structure functions for wind speed, zonal and meridional wind components are estimated from ASCAT retrievals. Furthermore, the new procedure includes a temporal interpolation of the retrievals based on the complex empirical orthogonal function (CEOF) approach, in order to enhance the sampling length of the scatterometer observations. The resulting daily wind fields involves the main known surface wind patterns as well as some variation modes associated with temporal and spatial moving features. The accuracy of the gridded winds was investigated through comparisons with moored buoy data in Bentamy et al. (2012) and resulted in rms differences for wind speed and direction are about 1.50 m.s⁻¹ and 20°.

2.3.3 ISAS

The In Situ Analysis System (ISAS), as described in Gaillard et al. (2016) is a data based reanalysis of temperature and salinity fields over the global ocean. It was initially designed to synthesize the temperature and salinity profiles collected by the Argo program. It has been later extended to accommodate all type of vertical profile as well as time series. ISAS gridded fields are entirely based on in-situ measurements. The methodology and configuration have been conceived to preserve as much as possible the data information content and resolution. ISAS is developed and run in a research laboratory (LOPS) in close collaboration with Coriolis, one of Argo Global Data Assembly Center and unique data provider for the Mercator operational oceanography system. At the moment the period covered starts in 2002 and only the upper 2000 m are considered. The gridded fields were produced over the global ocean 70°N-70°S on a 1/2° grid by the ISAS project with datasets downloaded from the Coriolis data center (for more details on ISAS see Gaillard et al. (2009)). In the Pi-MEP, the product in used is the INSITU_GLO_TS_OA_NRT_OBSERVATIONS_013_002_a v6.2 NRT derived at the Coriolis data center and provided by Copernicus (www.marine.copernicus.eu/documents/ PUM/CMEMS-INS-PUM-013-002-ab.pdf). The major contribution to the data set is from Argo array of profiling floats, reaching an approximate resolution of one profile every 10-days and every 3-degrees over the satellite SSS period (http://www.umr-lops.fr/SNO-Argo/Products/ ISAS-T-S-fields/); in this version SSS from ship of opportunity thermosalinographs are not used, so that we can consider SMOS SSS validation using these measurements independent of ISAS. The ISAS optimal interpolation involves a structure function modeled as the sum of two



Gaussian functions, each associated with specific time and space scales, resulting in a smoothing over typically 3 degrees. The smallest scale which can be retrieved with ISAS analysis is not smaller than 300–500 km (Kolodziejczyk et al. (2015)). For validation purpose, the ISAS monthly SSS fields at 5 m depth are collocated and compared with the satellite SSS products and included in the Pi-MEP Match-up files. In addition, the "percentage of variance" fields (PCTVAR) contained in the ISAS analyses provide information on the local variability of in situ SSS measurements within $1/2^{\circ}x1/2^{\circ}$ boxes.

2.3.4 World Ocean Atlas Climatology

The World Ocean Atlas 2013 version 2 (WOA13 V2) is a set of objectively analyzed (1° grid) climatological fields of in situ temperature, salinity and other variables provided at standard depth levels for annual, seasonal, and monthly compositing periods for the World Ocean. It also includes associated statistical fields of observed oceanographic profile data interpolated to standard depth levels on 5°, 1°, and 0.25° grids. We use these fields in complement to ISAS to characterize the climatological fields (annual mean and std) at the match-up pairs location and date.

2.4 Overview of the Match-ups generation method

The match-up production is basically a three steps process:

- 1. preparation of the input in situ and satellite data, and,
- 2. co-localization of satellite products with in situ SSS measurements.
- 3. co-localization of the in situ/satellite pair with auxiliary information.

In the following, we successively detail the approaches taken for these different steps.

2.4.1 In Situ/Satellite data filtering

The first step consist in filtering Surface drifters in situ dataset using the quality flags as described in 2.2 so that only valid salinity data remains in the produced match-ups.

For high-spatial resolution in situ SSS measurements such as the Thermo-SalinoGraph (TSG) SSS data from research vessels, Voluntary Observing Ships (VOS) or sailing ships, as well as SSS data from surface drifters, an additional spatial-filtering step is performed on the in situ data that will be in fine compared to the satellite SSS products. If R_{sat} is the spatial resolution of the satellite SSS product (L2 to L3-L4), we keep the in situ data at the original spatial resolution but we also estimate for all spatio-temporal samples a running median filtered SSS applied to all neighbouring in situ SSS data acquired within a distance of $R_{sat}/2$ from a given in situ acquisition. Both the original and the filtered data are finally stored in the MDB files.

Only for satellite L2 SSS data, a third step consist in filtering spurious data using the flags and associated recommendation as provided by the official data centers and described in 2.1.

2.4.2 In Situ/Satellite Co-localization

In this step, each SSS satellite acquisition is co-localized with the filtered in situ measurements. The method used for co-localization differ if the satellite SSS is a swath product (so-called Level 2-types) or a time-space composite product (so-called Level 3/level 4-types).



• For L2 SSS swath data:

If R_{sat} is the spatial resolution of the satellite swath SSS product, for each in situ data sample collected in the Pi-MEP database, the platform searches for all satellite SSS data found at grid nodes located within a radius of $R_{sat}/2$ from the in situ data location and acquired with a time-lag from the in situ measurement date that is less or equal than \pm 12 hours. If several satellite SSS samples are found to meet these criteria, the final satellite SSS match-up point is selected to be the closest in time from the in situ data measurement date. The final spatial and temporal lags between the in situ and satellite data are stored in the MDB files.

• For L3 and L4 composite SSS products:

If R_{sat} is the spatial resolution of the composite satellite SSS product and D the period over which the composite product was built (e.g., periods of 1, 7, 8, 9, 10, 18 days, 1 month, etc..) with central time to, for each in situ data sample collected in the Pi-MEP database during period D, the platform searches for all satellite SSS data of the composite product found at grid nodes located within a radius of $R_{sat}/2$ from the in situ data location. If several satellite SSS product samples are found to meet these criteria, the final satellite SSS match-up point is chosen to be the composite SSS with central time to which is the closest in time from the in situ data measurement date. The final spatial and temporal lags between the in situ and satellite data are stored in the MDB files.

2.4.3 MDB pair Co-localization with auxiliary data and complementary information

MDB data consist of satellite and in-situ SSS pair datasets but also of auxiliary geophysical parameters such as local and history of wind speed and rain rates, as well as various information (climatology, distance to coast, mixed layer depth, barrier layer thickness, etc) that can be derived from in situ data and which are included in the final match-up files. The collocation of auxiliary parameters and additional information is done for each filtered in-situ SSS measurement contained in the match-up files as follows:

If t_{insitu} is the time/date at which the in situ measurement is performed, we collect:

- The ASCAT wind speed product of the same day than t_{insitu} found at the ASCAT $1/4^{\circ}$ grid node with closest distance from the in situ data location and the time series of the ASCAT wind speed at the same node for the 10 days prior the in situ measurement day.
- If the in situ data is located within the 60°N-60°S band, we select the CMORPH 3-hourly product the closest in time from tin situ and found at the CMORPH 1/4° grid node with closest distance from the in situ data location. We then store the time series of the CMORPH rain rate at the same node for the 10 days prior the in situ measurement time.

For the given month/year of the in situ data, we select the ISAS and WOA fields for the same month (and same year for ISAS fields) and take the SSS analysis (monthly mean, std) found at the closest grid node from the in situ measurement.

The distance from the in situ SSS data location to the nearest coast is evaluated and provided in km. We use a distance-to-coast map at $1/4^{\circ}$ resolution where small islands have been removed.

When vertical profiles of salinity (S) and temperature (T) are made available from the in situ measurements used to build the match-up (Argo or sea mammals), the following variables are included into each satellite/in situ match-up file:

1. The vertical distribution of pressure at which the profile were measured,



- 2. The vertical S(z) and T(z) profiles,
- 3. The vertical potential density anomaly profile $\sigma_0(z)$,
- 4. The Mixed Layer Depth (MLD). The MLD is defined here as the depth where the potential density has increased from the reference depth (10 meter) by a threshold equivalent to 0.2°C decrease in temperature at constant salinity: $\sigma_0 = \sigma_{010m} + \Delta \sigma_0$ with $\Delta \sigma_0 = \sigma_0(\theta_{10m} 0.2, S_{10m}) \sigma_0(\theta_{10m}, S_{10m})$ where θ_{10m} and S_{10m} are the temperature and salinity at the reference depth (i.e. 10 m) (de Boyer Montégut et al. (2004), de Boyer Montégut et al. (2007)).
- 5. The Top of the Thermocline Depth (TTD) is defined as the depth at which temperature decreases from its 10 m value by 0.2°C.
- 6. The Barrier Layer if present, is defined as the intermediate layer between the top of the thermocline and the bottom of the density mixed-layer and its thickness (BLT) is defined as the difference between the MLD and the TTD.
- 7. The vertical profile of the buoyancy frequency $N^2(z)$

The resulting match-ups files are serialized as NetCDF-4 files whose structure depends on the origin of the in-situ data they contain.

2.4.4 Content of the Match-Up NetCDF files

```
netcdf pimep-mdb_aquarius-l3-or-v5-7dr-rain-mask_drifter_TIMEID_v01 {
dimensions:
    TIME\_SAT = UNLIMITED; // (1 currently)
    TIME_DRIFTER = 199;
    N_DAYS_WIND = 10;
    N_3H_RAIN = 80;
  variables:
float DATE_DRIFTER(TIME_DRIFTER);
DATE_DRIFTER:long_name = "Date of drifter";
    DATE_DRIFTER:units = "days since 1990-01-01 00:00:00";
    DATE_DRIFTER:standard_name = "time";
    DATE_DRIFTER:_FillValue = -999.f;
float LATITUDE_DRIFTER(TIME_DRIFTER);
    LATITUDE_DRIFTER:long_name = "Latitude of drifter";
    LATITUDE_DRIFTER:units = "degrees_north";
    LATITUDE_DRIFTER:valid_min = -90.;
    LATITUDE_DRIFTER:valid_max = 90.;
    LATITUDE_DRIFTER:standard_name = "latitude";
    LATITUDE_DRIFTER:_FillValue = -999.f;
float LONGITUDE_DRIFTER(TIME_DRIFTER):
    LONGITUDE_DRIFTER:long_name = "Longitude of drifter";
    LONGITUDE\_DRIFTER: units = "degrees\_east" \; ;
    LONGITUDE_DRIFTER: valid_min = -180.;
    LONGITUDE_DRIFTER:valid_max = 180.;
    LONGITUDE_DRIFTER:standard_name = "longitude";
```



```
LONGITUDE_DRIFTER:_FillValue = -999.f;
float SSS_DRIFTER(TIME_DRIFTER) ;
    SSS_DRIFTER:long_name = "Drifter SSS";
    SSS_DRIFTER:units = "1";
    SSS_DRIFTER:salinity_scale = "Practical Salinity Scale(PSS-78)";
    SSS_DRIFTER:standard_name = "sea_water_salinity";
    SSS_DRIFTER:_FillValue = -999.f;
float SST_DRIFTER(TIME_DRIFTER) ;
    SST_DRIFTER:long_name = "Drifter SST";
    SST_DRIFTER:units = "degree Celsius";
    SST_DRIFTER:standard_name = "sea_water_temperature";
    SST_DRIFTER:_FillValue = -999.f;
float SSS_DRIFTER_FILTERED(TIME_DRIFTER);
    SSS_DRIFTER_FILTERED:long_name = "Drifter SSS median filtered at satellite spatial
resolution";
    SSS_DRIFTER_FILTERED:units = "1";
    SSS_DRIFTER_FILTERED:salinity_scale = "Practical Salinity Scale(PSS-78)";
    SSS_DRIFTER_FILTERED:standard_name = "sea_water_salinity";
    SSS_DRIFTER_FILTERED:_FillValue = -999.f;
float SST_DRIFTER_FILTERED(TIME_DRIFTER);
    SST_DRIFTER_FILTERED:long_name = "Drifter SST median filtered at satellite spatial
resolution";
    SST_DRIFTER_FILTERED:units = "degree Celsius";
    SST_DRIFTER_FILTERED:standard_name = "sea_water_temperature";
    SST_DRIFTER_FILTERED:_FillValue = -999.f;
float DISTANCE_TO_COAST_DRIFTER(TIME_DRIFTER) ;
    DISTANCE_TO_COAST_DRIFTER:long_name = "Distance to coasts at drifter location"
;
    DISTANCE_TO_COAST_DRIFTER:units = "km";
    DISTANCE_TO_COAST_DRIFTER:_FillValue = -999.f;
float PLATFORM_NUMBER_DRIFTER(TIME_DRIFTER) ;
    PLATFORM_NUMBER_DRIFTER:long_name = "drifter unique identifier";
    \label{eq:platform_number_def} PLATFORM\_NUMBER\_DRIFTER: conventions = "WMO float identifier: A9IIIII";
    PLATFORM_NUMBER_DRIFTER:units = "1":
    PLATFORM_NUMBER_DRIFTER:_FillValue = -999.f;
float DATE_Satellite_product(TIME_Sat);
    DATE_Satellite_product:long_name = "Central time of satellite SSS file";
    DATE_Satellite_product:units = "days since 1990-01-01 00:00:00";
    DATE_Satellite_product:standard_name = "time";
float LATITUDE_Satellite_product(TIME_DRIFTER) ;
    LATITUDE_Satellite_product:long_name = "Satellite product latitude at drifter location"
    LATITUDE_Satellite_product:units = "degrees_north";
    LATITUDE_Satellite_product:valid_min = -90.;
    LATITUDE_Satellite_product:valid_max = 90.;
    LATITUDE\_Satellite\_product:standard\_name = "latitude" \ ;
    LATITUDE_Satellite_product:_FillValue = -999.f;
float LONGITUDE_Satellite_product(TIME_DRIFTER);
    LONGITUDE_Satellite_product:long_name = "Satellite product longitude at drifter loca-
```



```
tion";
        LONGITUDE_Satellite_product:units = "degrees_east";
        LONGITUDE_Satellite_product:valid_min = -180.;
        LONGITUDE_Satellite_product:valid_max = 180.;
        LONGITUDE\_Satellite\_product: standard\_name = "longitude" \ ;
        LONGITUDE_Satellite_product:_FillValue = -999.f;
float SSS_Satellite_product(TIME_DRIFTER) ;
        SSS_Satellite_product:long_name = "Satellite product SSS at drifter location";
        SSS_Satellite_product:units = "1";
        SSS_Satellite_product:salinity_scale = "Practical Salinity Scale(PSS-78)";
        SSS_Satellite_product:standard_name = "sea_surface_salinity";
        SSS\_Satellite\_product:\_FillValue = -999.f;
float SST_Satellite_product(TIME_DRIFTER) ;
        SST_Satellite_product:long_name = "Satellite product SST at drifter location";
        SST_Satellite_product:units = "degree Celsius";
        SST_Satellite_product:standard_name = "sea_surface_temperature";
        SST\_Satellite\_product:\_FillValue = -999.f;
float Spatial_lags(TIME_DRIFTER) ;
         Spatial_lags:long_name = "Spatial lag between drifter location and satellite SSS product
pixel center";
        Spatial_lags:units = "km";
        Spatial_lags:_FillValue = -999.f;
float Time_lags(TIME_DRIFTER);
        Time_lags:long_name = "Temporal lag between drifter time and satellite SSS product cen-
tral time";
        Time_{lags:units} = "days";
        Time_lags:_FillValue = -999.f;
float ROSSBY_RADIUS_at_DRIFTER(TIME_DRIFTER) ;
         ROSSBY_RADIUS_at_DRIFTER:long_name = "Baroclinic Rossby radius of deformation
(Chelton et al., 1998) at drifter location";
        ROSSBY_RADIUS_at_DRIFTER:units = "km";
        ROSSBY_RADIUS_at_DRIFTER:_FillValue = -999.f;
float Ascat_daily_wind_at_DRIFTER(TIME_DRIFTER);
         Ascat_daily_wind_at_DRIFTER:long_name = "Daily Ascat wind speed module at drifter
location";
        Ascat_daily\_wind_at_DRIFTER:units = "m/s";
        Ascat_daily_wind_at_DRIFTER:_FillValue = -999.f;
float CMORPH_3h_Rain_Rate_at_DRIFTER(TIME_DRIFTER);
        {\it CMORPH\_3h\_Rain\_Rate\_at\_DRIFTER:} long\_name = "3-hourly CMORPH rain rate at drifter and all the content of 
location";
        CMORPH_3h_Rain_Rate_at_DRIFTER:units = "mm/3h";
        CMORPH_3h_Rain_Rate_at_DRIFTER:_FillValue = -999.f :
float Ascat_10_prior_days_wind_at_DRIFTER(TIME_DRIFTER, N_DAYS_WIND);
         Ascat_10_prior_days_wind_at_DRIFTER:long_name = "Prior 10 days time series of Ascat
wind speed module at drifter location";
        Ascat_10_prior_days_wind_at_DRIFTER:units = "m/s";
        Ascat_10_prior_days_wind_at_DRIFTER:_FillValue = -999.f;
float CMORPH_10_prior_days_Rain_Rate_at_DRIFTER(TIME_DRIFTER, N_3H_RAIN);
        CMORPH_10_prior_days_Rain_Rate_at_DRIFTER:long_name = "Prior 10 days times series
```



```
of 3-hourly CMORPH Rain Rate at drifter location";
    CMORPH_10_prior_days_Rain_Rate_at_DRIFTER:units = "mm/3h";
    CMORPH_10_prior_days_Rain_Rate_at_DRIFTER:_FillValue = -999.f;
float SSS_ISAS_at_DRIFTER(TIME_DRIFTER);
    SSS_ISAS_at_DRIFTER:long_name = "ISAS SSS (5m depth) at drifter location";
    SSS\_ISAS\_at\_DRIFTER:units = "1";
    SSS_ISAS_at_DRIFTER:salinity_scale = "Practical Salinity Scale(PSS-78)";
    SSS_ISAS_at_DRIFTER:standard_name = "sea_water_salinity";
    SSS_ISAS_at_DRIFTER:_FillValue = -999.f;
float SSS_PCTVAR_ISAS_at_DRIFTER(TIME_DRIFTER);
    SSS_PCTVAR_ISAS_at_DRIFTER:long_name = "Error on ISAS SSS (5m depth) at drifter
location (% variance)";
    SSS\_PCTVAR\_ISAS\_at\_DRIFTER:units = "%";
    SSS\_PCTVAR\_ISAS\_at\_DRIFTER:\_FillValue = -999.f;
float SSS_WOA13_at_DRIFTER(TIME_DRIFTER);
    SSS_WOA13_at_DRIFTER:long_name = "WOA 2013 (DECAV-1deg) SSS (0m depth) at
drifter location";
    SSS_WOA13_at_DRIFTER:units = "1";
    SSS_WOA13_at_DRIFTER:salinity_scale = "Practical Salinity Scale(PSS-78)";
    SSS_WOA13_at_DRIFTER:standard_name = "sea_surface_salinity";
    SSS_WOA13_at_DRIFTER:_FillValue = -999.f;
float SSS_STD_WOA13_at_DRIFTER(TIME_DRIFTER);
    SSS_STD_WOA13_at_DRIFTER:long_name = "WOA 2013 (DECAV-1deg) SSS STD (0m
depth) at drifter location ";
    SSS_STD_WOA13_at_DRIFTER:units = "1";
    SSS\_STD\_WOA13\_at\_DRIFTER:\_FillValue = -999.f;
float SSS_ISAS15_at_DRIFTER(N_prof);
    SSS_ISAS15_at_DRIFTER:long_name = "Monthly ISAS-15 SSS (5m depth) at drifter loca-
tion";
    SSS_ISAS15_at_DRIFTER:units = "1";
    SSS_ISAS15_at_DRIFTER:salinity_scale = "Practical Salinity Scale (PSS-78)";
    SSS_ISAS15_at_DRIFTER:standard_name = "sea_water_salinity";
    SSS_ISAS15_at_DRIFTER:_FillValue = -999.f;
float SSS_PCTVAR_ISAS15_at_DRIFTER(N_prof) :
     SSS\_PCTVAR\_ISAS15\_at\_DRIFTER:long\_name = "Error on monthly ISAS-15 SSS (5m)
depth) at drifter location (% variance)";
    SSS_PCTVAR_ISAS15_at_DRIFTER:units = "%";
    SSS_PCTVAR_ISAS15_at_DRIFTER:_FillValue = -999.f;
float SSS_WOA18_at_DRIFTER(N_prof);
     SSS_WOA18_at_DRIFTER:long_name = "Monthly WOA 2018 (DECAV-1deg) SSS (0m
depth) at drifter location";
    SSS_WOA18_at_DRIFTER:units = "1":
    SSS\_WOA18\_at\_DRIFTER: salinity\_scale = "Practical Salinity Scale (PSS-78)" \ ;
    SSS_WOA18_at_DRIFTER:standard_name = "sea_surface_salinity";
    SSS_WOA18_at_DRIFTER:_FillValue = -999.f;
float SSS_STD_WOA18_at_DRIFTER(N_prof);
     SSS_STD_WOA18_at_DRIFTER:long_name = "Monthly WOA 2018 (DECAV-1deg) SSS
STD (0m depth) at drifter location ";
    SSS_STD_WOA18_at_DRIFTER:units = "1";
```



```
SSS\_STD\_WOA18\_at\_DRIFTER:\_FillValue = -999.f;
float SEA_ICE_CONCENTRATION_at_DRIFTER(N_prof);
        SEA_ICE_CONCENTRATION_at_DRIFTER:long_name = "Daily sea ice area fraction (EU-
METSAT OSI-SAF OSI-450) at drifter location (%)";
        SEA_ICE_CONCENTRATION_at_DRIFTER:units = "1";
        SEA_ICE_CONCENTRATION_at_DRIFTER:standard_name = "sea_ice_area_fraction";
        SEA_ICE_CONCENTRATION_at_DRIFTER:_FillValue = -999.f;
float CCMP_6h_Wind_Speed_at_DRIFTER(N_prof) ;
        CCMP_6h_Wind_Speed_at_DRIFTER:long_name = "6-hourly CCMP wind speed at drifter
location":
        CCMP_6h_Wind_Speed_at_DRIFTER:units = "m s-1";
        \label{lem:comp_obj} \begin{split} & CCMP\_6h\_Wind\_Speed\_at\_DRIFTER:standard\_name = "wind\_speed" \ ; \end{split}
        CCMP_6h_Wind_Speed_at_DRIFTER:_FillValue = -999.f;
float CCMP_10_prior_days_Wind_Speed_at_DRIFTER(N_prof, N_DAYS_WIND_CCMP);
        CCMP_10_prior_days_Wind_Speed_at_DRIFTER:long_name = "Prior 10 days time series of
CCMP wind speed at drifter location";
        CCMP_10_prior_days_Wind_Speed_at_DRIFTER:units = "m s-1";
        CCMP_10_prior_days_Wind_Speed_at_DRIFTER:standard_name = "wind_speed";
        CCMP_10_prior_days_Wind_Speed_at_DRIFTER:_FillValue = -999.f;
float CDM_GLOBCOLOUR_at_DRIFTER(N_prof);
        {\tt CDM\_GLOBCOLOUR\_at\_DRIFTER:long\_name} = "8-day\ Coloured\ dissolved\ and\ detrital
organic materials - mean of the binned pixels at drifter location";
        CDM_GLOBCOLOUR_at_DRIFTER:units = "m-1";
        CDM\_GLOBCOLOUR\_at\_DRIFTER: standard\_name = "volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_coefficient\_of\_radiative\_flux\_in\_sea\_volume\_absorption\_absorption\_coefficient\_of\_radiative\_absorption\_coefficient\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorption\_absorptio
        CDM_GLOBCOLOUR_at_DRIFTER:_FillValue = -999.f;
float CHL1_GLOBCOLOUR_at_DRIFTER(N_prof);
        CHL1_GLOBCOLOUR_at_DRIFTER:long_name = "8-day Chlorophyll concentration - mean
of the binned pixels at drifter location";
        CHL1_GLOBCOLOUR_at_DRIFTER:units = "mg m-3";
        CHL1\_GLOBCOLOUR\_at\_DRIFTER: standard\_name = "mass\_concentration\_of\_chlorophyll\_a\_in\_sea\_water"
        CHL1_GLOBCOLOUR_at_DRIFTER:_FillValue = -999.f;
float EVAPORATION_OAFLUX_at_DRIFTER(N_prof) :
        EVAPORATION_OAFLUX_at_DRIFTER:long_name = "Daily mean evaporation rate (OAFlux)
at drifter location";
        EVAPORATION_OAFLUX_at_DRIFTER:units = "cm year-1";
        EVAPORATION_OAFLUX_at_DRIFTER:_FillValue = -999.f;
float SSS_SCRIPPS_at_DRIFTER(N_prof) ;
        SSS_SCRIPPS_at_DRIFTER:long_name = "Argo gridded monthly mean SSS (0m depth)
from SCRIPPS (Roemmich-Gilson) at drifter location";
        SSS_SCRIPPS_at_DRIFTER:units = "1";
        SSS_SCRIPPS_at_DRIFTER:salinity_scale = "Practical Salinity Scale (PSS-78)";
        SSS\_SCRIPPS\_at\_DRIFTER: standard\_name = "sea\_water\_salinity" \ ;
        SSS_SCRIPPS_at_DRIFTER:_FillValue = -999.f;
float SSS_IPRC_at_DRIFTER(N_prof);
        SSS_IPRC_at_DRIFTER:long_name = "Argo gridded monthly mean SSS (0m depth) from
IPRC at drifter location";
        SSS\_IPRC\_at\_DRIFTER:units = "1";
```



```
SSS_IPRC_at_DRIFTER:salinity_scale = "Practical Salinity Scale (PSS-78)";
    SSS_IPRC_at_DRIFTER:standard_name = "sea_water_salinity";
    SSS\_IPRC\_at\_DRIFTER:\_FillValue = -999.f;
float SST_AVHRR_at_DRIFTER(N_prof);
    SST_AVHRR_at_DRIFTER:long_name = "Daily OI AVHRR-only v2 SST (Reynolds et al.,
2007) at drifter location";
    SST_AVHRR_at_DRIFTER:units = "degree Celsius";
    SST\_AVHRR\_at\_DRIFTER: standard\_name = "sea\_water\_temperature" \ ;
    SST_AVHRR_at_DRIFTER:_FillValue = -999.f;
float U_EKMAN_GLOBCURRENT_at_DRIFTER(N_prof);
    U_EKMAN_GLOBCURRENT_at_DRIFTER:long_name = "15m depth Ekman current ve-
locity: zonal component at drifter location";
    U_EKMAN_GLOBCURRENT_at_DRIFTER:units = "m s-1";
    U_EKMAN_GLOBCURRENT_at_DRIFTER:_FillValue = -999.f;
float V_EKMAN_GLOBCURRENT_at_DRIFTER(N_prof);
    V_EKMAN_GLOBCURRENT_at_DRIFTER:long_name = "15m depth Ekman current ve-
locity: meridian component at drifter location";
    V_EKMAN_GLOBCURRENT_at_DRIFTER:units = "m s-1";
    V_EKMAN_GLOBCURRENT_at_DRIFTER:_FillValue = -999.f;
float U_GEOSTROPHIC_GLOBCURRENT_at_DRIFTER(N_prof);
    U_GEOSTROPHIC_GLOBCURRENT_at_DRIFTER:long_name = "Absolute geostrophic
velocity: zonal component at drifter location";
    U_GEOSTROPHIC_GLOBCURRENT_at_DRIFTER:units = "m s-1";
    U_GEOSTROPHIC_GLOBCURRENT_at_DRIFTER:_FillValue = -999.f;
float V_GEOSTROPHIC_GLOBCURRENT_at_DRIFTER(N_prof);
    V_GEOSTROPHIC_GLOBCURRENT_at_DRIFTER:long_name = "Absolute geostrophic
velocity: meridian component at drifter location";
    V_GEOSTROPHIC_GLOBCURRENT_at_DRIFTER:units = "m s-1";
    V_GEOSTROPHIC_GLOBCURRENT_at_DRIFTER:_FillValue = -999.f;
   // global attributes:
    :Conventions = "CF-1.6";
    :title = "Surface drifters Match-Up Database";
    :Satellite_product_name = "Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG";
    :Satellite_product_spatial_resolution = "1 deg";
    :Satellite_product_temporal_resolution = "7 day running";
    :Satellite_product_filename = "v5/7day_running/SCI/2011/237/Q20112312011237.L3m_R7_SCI_V5.0.RAIN_MASK_SSS_ldeg.bz2"
    :Match-Up_spatial_window_radius_in_km = 55;
    :Match-Up_temporal_window_radius_in_days = 0.25;
    :start\_time = "20100114T000005Z";
    :stop\_time = "20100118T235026Z";
    :northernmost_latitude = 77.676f;
    :sourthenmost_latitude = -66.423f;
    :westernmost_longitude = -179.219f;
    :easternmost_longitude = 179.199f;
    :geospatial_lat_units = "degrees north";
    :geospatial_lat_resolution = "1 deg";
    :geospatial_lon_units = "degrees east";
```



```
:geospatial.lon_resolution = "1 deg";
:institution = "ESA-IFREMER-ODL";
:project_name = "SMOS Pilote Mission Exploitation Platfrom (Pi-MEP) for salinity";
:project_url = "https://pimep-project.odl.bzh";
:license = "Pi-MEP data use is free and open";
:product_version = "1.0";
:keywords = "Oceans > Ocean Salinity > Sea Surface Salinity";
:acknowledgment = "Please acknowledge the use of these data with the following statement:
These data were provided by SMOS Pilote Mission Exploitation Platfrom (Pi-MEP) for salinity";
:source = "v5/7day_running/SCI/2011/237/Q20112312011237.L3m_R7_SCI_V5.0.RAIN_MASK_SSS_1deg.bz2";
:In_situ_data_source = "https://skyros.locean-ipsl.upmc.fr/~smos/drifters/valid/";
:references = "https://pimep-project.odl.bzh";
:history = "Processed on 2018-04-18 using MDB_generator";
:date_created = "2018-04-18 17:09:30";
}
```

2.5 MDB characteristics for the particular in situ/satellite pairs

2.5.1 Number of paired SSS data as a function of time and distance to coast




Figure 1: Number of match-ups between Surface drifters and Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG SSS as a function of time (a) and as function of the distance to coast (b) over the Tropical Pacific Ocean Pi-MEP region and for the full satellite product period.



2.5.2 Histograms of the SSS match-ups




Figure 2: Histograms of SSS from Surface drifters (a) and Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG (b) considering all match-up pairs per bins of 0.1 over the Tropical Pacific Ocean Pi-MEP region and for the full satellite product period.

2.5.3 Spatial Distribution of Match-ups




Figure 3: Number of SSS match-ups between Surface drifters SSS and the Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG SSS product for the Tropical Pacific Ocean Pi-MEP region over 1°x1° boxes and for the full satellite product period.

2.5.4 Histograms of the spatial and temporal lags of the match-ups pairs




Figure 4: Histograms of the spatial (a) and temporal (b) lags between the time of the Surface drifters measurements and the date of the corresponding Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG SSS product.



3 MDB file Analyses

3.1 Spatial Maps of the Temporal mean and Std of in situ and satellite SSS and of the difference (Δ SSS)

In Figure 5, we show maps of temporal mean (left) and standard deviation (right) of the Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG satellite SSS product (top) and of the Surface drifters in situ dataset at the collected Pi-MEP match-up pairs. The temporal mean and std are gridded over the full satellite product period and over spatial boxes of size 1° x1°.

At the bottom of Figure 5, the temporal mean (left) and standard deviation (right) of the differences between the satellite SSS product and in situ data found at match-up pairs, namely Δ SSS(Satellite -Surface drifters), is also gridded over the full satellite product period and over spatial boxes of size 1°x1°.




Figure 5: Temporal mean (left) and Std (right) of SSS from Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG (top), Surface drifters (middle), and of Δ SSS (Satellite - Surface drifters). Only match-up pairs are used to generate these maps.

3.2 Time series of the monthly averaged mean and Std of in situ and satellite SSS and of the (Δ SSS)

In the top panel of Figure 6, we show the time series of the monthly averaged SSS estimated over the full Tropical Pacific Ocean Pi-MEP region for both Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG satellite SSS product (in black) and the Surface drifters in situ dataset (in blue) at the collected Pi-MEP match-up pairs.

In the middle panel of Figure 6, we show the time series of the monthly averaged Δ SSS (Satellite - Surface drifters) for the collected Pi-MEP match-up pairs and estimated over the full Tropical Pacific Ocean Pi-MEP region.

In the bottom panel of Figure 6, we show the time series of the monthly averaged standard deviation of the Δ SSS (Satellite - Surface drifters) for the collected Pi-MEP match-up pairs and



estimated over the full Tropical Pacific Ocean Pi-MEP region.




Figure 6: Time series of the monthly averaged mean SSS (top), mean Δ SSS (Satellite - Surface drifters) and Std of Δ SSS (Satellite - Surface drifters) over the Tropical Pacific Ocean Pi-MEP region considering all match-ups collected by the Pi-MEP platform.

3.3 Zonally-averaged Time-mean and temporal Std of in situ and satellite SSS and of the Δ SSS

In Figure 7 left panel, we show the zonally averaged time-mean SSS estimated at the collected Pi-MEP match-up pairs for both Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG satellite SSS product (in black) and the Surface drifters in situ dataset (in blue). The time mean is evaluated over the full satellite SSS product period.

In the right panel of Figure 7, we show the zonally averaged time-mean Δ SSS (Satellite -Surface drifters) for all the collected Pi-MEP match-up pairs estimated over the full satellite product period.






Figure 7: Left panel: Zonally averaged time mean SSS from Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG (black) and from Surface drifters (blue). Right panel: zonally averaged time-mean Δ SSS (Satellite - Surface drifters) for all the collected Pi-MEP match-up pairs estimated over the full satellite product period.



3.4 Scatterplots of satellite vs in situ SSS by latitudinal bands




Figure 8: Contour maps of the concentration of Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG SSS (y-axis) versus Surface drifters SSS (x-axis) at match-up pairs for different latitude bands. For each plot, the red line shows x=y. The black thin and dashed lines indicate a linear fit through the data cloud and the $\pm 95\%$ confidence levels, respectively. The number match-up pairs n, the slope and R^2 coefficient of the linear fit, the root mean square (RMS) and the mean bias between satellite and in situ data are indicated for each latitude band in each plots.



3.5 Time series of the monthly averaged mean and Std of the ΔSSS sorted by latitudinal bands




Figure 9: Monthly-average mean (red curves) Δ SSS (Satellite - Surface drifters) and ± 1 Std (black vertical thick bars) as function of time for all the collected Pi-MEP match-up pairs estimated over the Tropical Pacific Ocean Pi-MEP region and for the full satellite product period are shown for different latitude bands: (a) Latitude band 80°S-80°N, (b) latitude band 20°S-20°N, (c) Mid Latitude bands 40°S-20°S and 20°N-40°N and (d) Latitude bands 60°S-40°S and 40°N-60°N.

3.6 Δ SSS sorted as function of geophysical parameters

In Figure 10, we classify the match-up differences Δ SSS (Satellite - in situ) between Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG and Surface drifters SSS as function of the geophysical conditions at match-up points. The mean and std of Δ SSS (Satellite - Surface drifters) is thus evaluated as function of the

- in situ SSS values per bins of width 0.2,
- \bullet in situ SST values per bins of width 1°C,
- ASCAT daily wind values per bins of width 1 m/s,
- CMORPH 3-hourly rain rates per bins of width 1 mm/h, and,
- distance to coasts per bins of width 50 km.






Figure 10: Δ SSS (Satellite - Surface drifters) sorted as function of Surface drifters SSS values a), Surface drifters SST b), ASCAT Wind speed c), CMORPH rain rate d) and distance to coast (e). In all plots the mean and Std of Δ SSS for each bin is indicated by the red curves and black vertical thick bars (± 1 Std)

3.7 Δ SSS maps and statistics for different geophysical conditions

In Figures 11 and 12, we focus on sub-datasets of the match-up differences Δ SSS (Satellite - in situ) between Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG and Surface drifters for the following specific geophysical conditions:

- C1:if the local value at in situ location of estimated rain rate is zero, mean daily wind is in the range [3, 12] m/s, the SST is > 5°C and distance to coast is > 800 km.
- C2:if the local value at in situ location of estimated rain rate is zero, mean daily wind is



in the range [3, 12] m/s.

- C3:if the local value at in situ location of estimated rain rate is high (ie. > 1 mm/h) and mean daily wind is low (ie. < 4 m/s).
- C5:if the in situ data is located where the climatological SSS standard deviation is low (ie. above < 0.2).
- C6:if the in situ data is located where the climatological SSS standard deviation is high (ie. above > 0.2).

For each of these conditions, the temporal mean (gridded over spatial boxes of size $1^{\circ}x1^{\circ}$) and the histogram of the difference ΔSSS (Satellite - in situ) are presented.




Figure 11: Temporal mean gridded over spatial boxes of size 1°x1° of Δ SSS (Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG - Surface drifters) for 5 different subdatasets corresponding to:RR=0 mm/h, 3< U_{10} <12 m/s, SST>5°C, distance to coast > 800 km (a), RR=0 mm/h, 3< U_{10} <12 m/s (b), RR>1mm/h and U_{10} <4m/s (c), WOA2013 SSS Std<0.2 (d), WOA2013 SSS Std>0.2 (e).






Figure 12: Normalized histogram of Δ SSS (Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG - Surface drifters) for 5 different subdatasets corresponding to: RR=0 mm/h, $3 < U_{10} < 12$ m/s, SST>5°C, distance to coast > 800 km (a), RR=0 mm/h, $3 < U_{10} < 12$ m/s (b), RR>1mm/h and $U_{10} < 4$ m/s (c), WOA2013 SSS Std<0.2 (d), WOA2013 SSS Std>0.2 (e).

4 Summary

Table 1 shows the mean, median, standard deviation (Std), root mean square (RMS), interquartile range (IQR), correlation coefficient ($\rm r^2$) and robust standard deviation (Std*) of the matchup differences Δ SSS (Satellite - in situ) between Aquarius-L3-OR-V5-7DAY-RUNNING-RAIN-MASK-1DEG and Surface drifters derived over the Tropical Pacific Ocean Pi-MEP region and for the full satellite product period and for the following conditions:

- all: All the match-up pairs satellite/in situ SSS values are used to derive the statistics
- C1: only pairs where RR=0 mm/h, $3 < U_{10} < 12$ m/s, SST>5°C, distance to coast > 800 km
- \bullet C2: only pairs where RR=0 mm/h, 3 < U_{10} <12 m/s
- C3: only pairs where RR>1mm/h and U_{10} <4m/s
- C5: only pairs where WOA2013 SSS Std<0.2
- C6: only pairs at WOA2013 SSS Std>0.2
- C7a: only pairs with a distance to coast < 150 km.
- C7b: only pairs with a distance to coast in the range [150, 800] km.
- C7c: only pairs with a distance to coast > 800 km.
- C8a: only pairs where SST is $< 5^{\circ}$ C.
- C8b: only pairs where SST is in the range [5, 15]°C.



- C8c: only pairs where SST is > 15°C.
- C9a: only pairs where SSS is < 33.
- C9b: only pairs where SSS is in the range [33, 37].
- C9c: only pairs where SSS is > 37.

Table 1: Statistics of Δ SSS (Satellite - Surface drifters)

Condition	#	Median	Mean	Std	RMS	IQR	\mathbf{r}^2	\mathbf{Std}^{\star}
all	39128	-0.03	0.00	0.21	0.21	0.24	0.73	0.17
C1	344	0.08	0.04	0.11	0.11	0.14	0.96	0.09
C2	27444	-0.04	-0.02	0.18	0.18	0.21	0.75	0.16
C3	351	0.12	0.22	0.40	0.46	0.49	0.41	0.36
C5	952	0.19	0.22	0.22	0.31	0.31	0.83	0.24
C6	38121	-0.03	0.00	0.21	0.21	0.23	0.72	0.17
C7a	4805	0.18	0.22	0.32	0.39	0.35	0.53	0.26
C7b	33933	-0.04	-0.03	0.17	0.17	0.21	0.74	0.16
C7c	390	0.07	0.04	0.10	0.11	0.14	0.97	0.10
C8a	0	NaN	NaN	NaN	NaN	NaN	NaN	NaN
C8b	0	NaN	NaN	NaN	NaN	NaN	NaN	NaN
C8c	39128	-0.03	0.00	0.21	0.21	0.24	0.73	0.17
C9a	88	0.56	0.93	0.73	1.18	1.34	0.97	0.64
C9b	39040	-0.03	0.00	0.20	0.20	0.24	0.68	0.17
C9c	0	NaN	NaN	NaN	NaN	NaN	NaN	NaN

For the same conditions, Table 2 presents statistics of Δ SSS (Satellite - ISAS). Only ISAS SSS values with PCTVAR<80% are used to derive the statistics.

Table 2: Statistics of Δ SSS (Satellite - ISAS)

Table 2. Statistics of ASSS (Saterite						10110)		
Condition	#	Median	Mean	Std	RMS	IQR	\mathbf{r}^2	\mathbf{Std}^{\star}
all	39128	0.03	0.03	0.16	0.17	0.22	0.77	0.17
C1	344	0.17	0.14	0.10	0.17	0.14	0.96	0.12
C2	27444	0.02	0.03	0.16	0.16	0.20	0.77	0.15
C3	351	-0.02	0.01	0.17	0.17	0.25	0.74	0.19
C5	952	0.18	0.14	0.11	0.18	0.15	0.94	0.14
C6	38121	0.02	0.03	0.16	0.17	0.21	0.76	0.16
C7a	4805	0.13	0.12	0.24	0.27	0.36	0.62	0.28
C7b	33933	0.01	0.02	0.15	0.15	0.20	0.77	0.15
C7c	390	0.17	0.14	0.10	0.17	0.14	0.97	0.12
C8a	0	NaN	NaN	NaN	NaN	NaN	NaN	NaN
C8b	0	NaN	NaN	NaN	NaN	NaN	NaN	NaN
C8c	39128	0.03	0.03	0.16	0.17	0.22	0.77	0.17
C9a	88	-0.02	0.02	0.14	0.15	0.32	0.99	0.08
C9b	39040	0.03	0.03	0.17	0.17	0.22	0.72	0.17
C9c	0	NaN	NaN	NaN	NaN	NaN	NaN	NaN

Numerical values can be downloaded as csv files for Table 1 and Table 2.



References

- Abderrahim Bentamy and Denis Croize Fillon. Gridded surface wind fields from Metop/ASCAT measurements. *Int. J. Remote Sens.*, 33(6):1729–1754, March 2012. ISSN 1366-5901. doi: 10.1080/01431161.2011.600348.
- Abderrahim Bentamy, Semyon A. Grodsky, James A. Carton, Denis Croizé-Fillon, and Bertrand Chapron. Matching ASCAT and QuikSCAT winds. *J. Geophys. Res.*, 117(C2), February 2012. ISSN 0148-0227. doi: 10.1029/2011JC007479. C02011.
- Jaqueline Boutin, Y. Chao, W. E. Asher, T. Delcroix, R. Drucker, K. Drushka, N. Kolodziejczyk, T. Lee, N. Reul, G. Reverdin, J. Schanze, A. Soloviev, L. Yu, J. Anderson, L. Brucker, E. Dinnat, A. S. Garcia, W. L. Jones, C. Maes, T. Meissner, W. Tang, N. Vinogradova, and B. Ward. Satellite and In Situ Salinity: Understanding Near-Surface Stratification and Sub-footprint Variability. Bull. Am. Meterol. Soc., 97(8):1391–1407, 2016. ISSN 1520-0477. doi: 10.1175/bams-d-15-00032.1.
- Clément de Boyer Montégut, Gurvan Madec, A. S. Fischer, A. Lazar, and D. Ludicone. Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology. *J. Geophys. Res.*, 109(C12):C12003, December 2004. ISSN 0148-0227. doi: 10.1029/2004jc002378.
- Clément de Boyer Montégut, Juliette Mignot, Alban Lazar, and Sophie Cravatte. Control of salinity on the mixed layer depth in the world ocean: 1. General description. *J. Geophys. Res.*, 112(C6):C06011, June 2007. ISSN 0148-0227. doi: 10.1029/2006jc003953.
- Ralph R. Ferraro. SSM/I derived global rainfall estimates for climatological applications. *J. Geophys. Res.*, 1021:16715–16736, 07 1997. doi: 10.1029/97JD01210.
- Ralph R. Ferraro, Fuzhong Weng, Norman C. Grody, and Limin Zhao. Precipitation characteristics over land from the NOAA-15 AMSU sensor. *Geophys. Res. Lett.*, 27(17):2669–2672, 2000. doi: 10.1029/2000GL011665.
- Fabienne Gaillard, E. Autret, V. Thierry, P. Galaup, C. Coatanoan, and T. Loubrieu. Quality Control of Large Argo Datasets. *J. Atmos. Oceanic Technol.*, 26(2):337–351, 2012/10/10 2009. doi: 10.1175/2008JTECHO552.1.
- Fabienne Gaillard, Thierry Reynaud, Virginie Thierry, Nicolas Kolodziejczyk, and Karina von Schuckmann. In Situ-Based Reanalysis of the Global Ocean Temperature and Salinity with ISAS: Variability of the Heat Content and Steric Height. *J. Clim.*, 29(4):1305–1323, February 2016. ISSN 1520-0442. doi: 10.1175/jcli-d-15-0028.1.
- Robert J. Joyce, John E. Janowiak, Phillip A. Arkin, and Pingping Xie. CMORPH: A Method that Produces Global Precipitation Estimates from Passive Microwave and Infrared Data at High Spatial and Temporal Resolution. *J. Hydrometeorol.*, 5(3):487–503, June 2004. ISSN 1525-7541. doi: 10.1175/1525-7541(2004)005/0487:camtpg)2.0.co;2.
- Nicolas Kolodziejczyk, Gilles Reverdin, and Alban Lazar. Interannual Variability of the Mixed Layer Winter Convection and Spice Injection in the Eastern Subtropical North Atlantic. *J. Phys. Oceanogr.*, 45(2):504–525, Feb 2015. ISSN 1520-0485. doi: 10.1175/jpo-d-14-0042.1.
- Christian Kummerow, Y. Hong, W. S. Olson, S. Yang, R. F. Adler, J. McCollum, R. Ferraro, G. Petty, D-B. Shin, and T. T. Wilheit. The Evolution of the Goddard Profiling Algorithm



(GPROF) for Rainfall Estimation from Passive Microwave Sensors. J. Appl. Meteorol., 40(11): 1801–1820, 2001. doi: $10.1175/1520-0450(2001)040\langle1801:TEOTGP\rangle2.0.CO;2.$